

MALLORY TRICHROME KIT

IVD *In vitro* diagnostic medical device



Three-reagent muscle and collagen fiber staining kit

INSTRUCTIONS FOR USE

REF Catalogue number: MT-100T (for 100 tests) MT-K-100 (3 x 100 ml)

Introduction

Mallory trichrome staining kit is used for treating the tested microscopic sample using three different stainings with differential counterstaining of two basic parts of the tissue (muscle and collagen fibers) in focus. By staining the sample with Fuchsin Acid acidic dye nuclei and muscles are stained red to pink. Phosphomolybdic acid molecules then push out Fuchsin Acid dye molecules from collagen and thus enables Aniline Blue to bind, resulting in collagen being stained contrast blue instead of red. Orange G (molecule of the lowest molar mass) stains erythrocytes.

Product description

- MALLORY TRICHROME KIT** - Three-reagent muscle and collagen fiber staining kit

The kit contains:	100 tests (MT-100T)	3 x 100 mL (MT-K-100)
Fuchsin Acid reagent	30 mL (FA-OT-30)	100 mL (FA-OT-100)
Phosphomolybdic acid, 1% solution	30 mL (FMK1-OT-30)	100 mL (FMK1-OT-100)
Orange G/Aniline Blue reagent	30 mL (OGA-OT-30)	100 mL (OGA-OT-100)

Other sections and reagents that may be used in staining:

- Fixatives such as BioGnost's neutral buffered formaldehyde solutions: Formaldehyde NB 4%, Formaldehyde NB 10%
- Dehydrating/rehydrating agent, such as BioGnost's alcohol solutions: Histanol 70, Histanol 80, Histanol 95 and Histanol 100
- Clearing agents, such as BioClear xylene or a substitute, such as BioClear New agent on the aliphatic hydrocarbons basis
- Infiltration and fitting agent, such as BioGnost's granulated paraffin BioWax Plus, BioWax 56/68, BioWax Blue, BioWax Micro.
- Covering agents for microscopic sections and mounting cover glass, such as BioGnost's BioMount, BioMount High, BioMount M, BioMount New, BioMount New Low, BioMount DPX, BioMount DPX High, BioMount DPX Low, BioMount DPX Low Eco, BioMount C, BioMount Aqua, Canada Balsam
- High-quality glass slides for use in histopathology and cytology, such as VitroGnost SUPER GRADE, VitroGnost COLOR or one of more than 30 models of BioGnost's VitroGnost glass slides
- VitroGnost cover glass, dimensions range from 18x18mm to 24x60mm
- BioGnost's immersion media, such as Immersion oil, Immersion oil, types A, C, FF, 37, or Immersion oil Tropical Grade

Preparing the histological sections for staining

- Fixate the sample (Formaldehyde NB 4%, Formaldehyde NB 10%), rinse with water and dehydrate through series of ascending alcohol solutions (Histanol 70, Histanol 80, Histanol 95 and Histanol 100).
- Clear the sample with intermedium; in xylene (BioClear) or in a xylene substitute (BioClear New).
- Infiltrate and fit the sample in paraffin (BioWax Plus, BioWax 56/58, BioWax Blue, BioWax Micro).
- Cut the paraffin block to 4-6 μ m slices and place them on a VitroGnost glass slide.

NOTE

Apply the reagent so it completely covers the section.

Sample staining procedure

a) using kit for 100 tests (MT-100T)

1.	Deparaffinize the section in xylene (BioClear) or in a xylene substitute (BioClear New)	3 exchanges, 2 min each
2.	Rehydrate using 100% alcohol (Histanol 100)	2 exchanges, 5 and 3 min
3.	Rehydrate using 95% alcohol (Histanol 95)	2 min
4.	Rehydrate in distilled (demi) water	2 min
5.	Stain with Fuchsin Acid reagent (add ≥ 5 drops)	30 seconds
6.	Rinse in distilled (demi) water	until the excessive dye is washed off of the section
7.	Add Phosphomolybdic acid, 1% solution (≥ 5 drops)	3 min
8.	Pour the reagent off the section without rinsing	
9.	Stain using Orange G/Aniline Blue reagent (add ≥ 5 drops)	4-6 minutes
10.	Dehydrate using 70% alcohol (Histanol 70)	5 dips
11.	Dehydrate using 95% alcohol (Histanol 95)	5 dips
12.	Dehydrate using 100% alcohol (Histanol 100)	2 min
13.	Clear the section in xylene (BioClear) or in a xylene substitute (BioClear New)	2 exchanges, 2 min each

Immediately after clearing apply an appropriate BioMount medium for covering/mounting on the section. If BioClear xylene was used, use one of BioGnost's mounting xylene-based media (BioMount, BioMount High, BioMount M, BioMount DPX, BioMount C, or universal BioMount New). If BioClear New xylene substitute was used, the appropriate covering agent is BioMount New. Cover the section with a VitroGnost cover glass.

b) using three-reagent 100 mL kit (MT-K-100)

Pour the reagents into glass staining jars (Coplin, Hellendahl or Schifferdecker), return to original bottles after staining. Close tightly. Filter the reagents if necessary.

1.	Deparaffinize the section in xylene (BioClear) or in a xylene substitute (BioClear New)	3 exchanges, 2 min each
2.	Rehydrate using 100% alcohol (Histanol 100)	2 exchanges, 5 and 3 min
3.	Rehydrate using 95% alcohol (Histanol 95)	2 min
4.	Rehydrate in distilled (demi) water	2 min
5.	Immerse into Fuchsin Acid reagent	30 seconds
6.	Rinse in distilled (demi) water	until the excessive dye is washed off of the section
7.	Immerse into Phosphomolybdic acid, 1% solution	3 min
8.	Remove excessive reagent from the section using filter paper (without rinsing in distilled water)	
9.	Immerse in Orange G/Aniline Blue reagent	4-6 minutes
10.	Dehydrate using 70% alcohol (Histanol 70)	5 dips
11.	Dehydrate using 95% alcohol (Histanol 95)	5 dips
12.	Dehydrate using 100% alcohol (Histanol 100)	2 min
13.	Clear the section in xylene (BioClear) or in a xylene substitute (BioClear New)	2 exchanges, 2 min each

Immediately after clearing apply an appropriate BioMount medium for covering/mounting on the section. If BioClear xylene was used, use one of BioGnost's mounting xylene-based media (BioMount, BioMount High, BioMount M, BioMount DPX, BioMount C, or universal BioMount New). If BioClear New xylene substitute was used, the appropriate covering agent is BioMount New. Cover the section with a VitroGnost cover glass.

Result

Muscle fibers, cytoplasm, nuclei - red to pink
Collagen - blue
Erythrocytes - orange to red

Note

Staining procedures are not standardized and they depend on standard operating procedures of individual laboratories and the experience of the personnel conducting the staining procedure. Intensity of staining depends on the period of immersion in the dye. Depending on personal requests and standard laboratory operating procedures, sample processing and staining can be carried out according to other protocols.

Preparing the sample and diagnostics

Use only appropriate instruments for collecting and preparing the samples. Process the samples with modern technology and mark them clearly. Follow the manufacturer's instructions for handling. In order to avoid mistakes, the staining procedure and diagnostics should only be conducted by authorized and qualified personnel. Use only microscope according to standards of the medical diagnostic laboratory.

Safety at work and environmental protection

Handle the product in accordance with safety at work and environmental protection guidelines. Used solutions and out of date solutions should be disposed of as special waste in accordance with national guidelines. Chemicals used in this procedure could pose danger to human health. Tested tissue specimens are potentially infectious. Necessary safety measures for protecting human health should be taken in accordance with good laboratory practice. Act in accordance with signs and warnings notices printed on the product's label, as well as in BioGnost's material safety data sheet.














Storing, stability and expiry date

Keep Mallory Trichrome kit in a tightly sealed original packaging at temperature of +15 to +25°C. Keep in dry places, do not freeze and avoid exposing to direct sunlight. Date of manufacture and expiry date are printed on the product's label.

References

1. Melis, M., Carpino, F., Di Tondo, U. (1989), Tecniche in anatomia patologica, Edi Ermes, Milano.
2. Prophet, E.B., Mills, B., Arrington, J., Sobin, L. (1968), Laboratory methods in histotechnology, McGraw Hill, Washington D.C.
3. Bancroft, J.D., Gamble, M. (2002), Theory and practice of Histological Techniques, Churchill Livingstone, New York.

MT-X, V6-EN6, November 24, 2021, IŠP/VR

 Refer to the supplied documentation	 Storage temperature range	 Number of tests in package	 Product code	 European Conformity
 Refer to supplied instructions	 Keep away from heat and sunlight	 Valid until	 Lot number	 Manufacturer
 For <i>in vitro</i> diagnostic use only	 Keep in dry place	 Caution - fragile		

 BIOGNOST Ltd.
Medjugorska 59
10040 Zagreb
CROATIA
www.biognost.com

