BIOGNOST®



Our global presence

We manufacture products for use in histopathology, hematology, cytology, and microbiology with a wide national and international distribution network present in over 70 countries.

BioGnost offers only the best quality products at competitive prices and has one of the widest ranges of medical products in South-Eastern Europe. Our experience and flexibility towards the client makes us a sought-after business partner, which has been recognized in the global market.



Our entrepreneurship began in 1990 with the founding of our family company. Our business growth began in 1998 with production of blood typing serums followed by Croatian Ministry of Health granting BioGnost the License for the manufacture of medical products. We are proud of one of our biggest successes - relocating to our new commercial and manufacturing building in 2012. The site enabled an enhanced work environment, including a carefully devised warehouse structure, segmented production facilities with increased manufacturing capacity, and additional R&D capabilities.

We continue to build on the strong momentum from last year to create high-quality products and provide excellent customer service.

strategic priorities

Our mission is to create high-quality products and provide excellent services, as well as introduce new or expanded production lines and sales ranges in accordance with market dynamics and demands.


Product and service excellence makes BioGnost a desirable regional and international business partner. The key to success is the ever-expanding network of business collaborations, as well as recognizable social responsibility. In the coming years we plan to further our market expansion, additionally invest in research and development, and to become competitive manufacturer in the fields of microbiological diagnostics and histopathology.

ISO 9001 ISO 13485 ISO 14001 BioGnost's entire business is in accordance with international ISO 9001, ISO 13485 and ISO 14001 standards. The company implemented ISO standards more than 15 years ago, and it is the first Croatian company to implement the ISO 13485 quality system for medical devices. Having fulfilled the requirements of these standards, the company is completely in accordance with national legal regulation and the European Union's Medical Device Directive.









CE IVD BioGnost's products bear the CE mark, a certificate that confirms that the products are in accordance with the European Union's standards. They are declared as *in vitro* diagnostic medical devices (IVD) classified as "Others" (according to the Directive 98/79/EC). No control of the Notified Body is required for this class; the company conducts compliance with valid regulations.

OEM manufacture

Besides products from our own sales ranges, we also offer OEM manufacture for your company. It includes a wide range of services, such as label design, product packaging, instructions for use and certificates of analysis, and products manufactured according to your recipes.



Diagnostic and clinical products

- 8 Marking systems
- 10 Formalin fixatives
- 11 Other fixatives
- 13 Decalcifying solutions
- 14 Dehydrating and rehydrating agents
- 15 Clearing and deparaffinization agents
- 16 Mounting media
- 20 Immersion oils
- 21 Cleaning solutions for microscope lens and slides
- 22 Infiltration, embedding and demolding media

Standard and special diagnostic stains


- 26 Histology stains
- 30 Cytology stains
- 32 Hematology stains
- 36 Bacteriology stains
- 41 Fungal stains
- 42 Special stains for histology and cytology
- 58 Powder dyes and stains for use in microscopy


Laboratory diagnostics

- 64 VitroGnost slides for microscopy
- 6/ Sample transport containers
- 68 Cassettes for histopathology
- 70 Pads and molds for histology
- 70 Equipment and accessories
- 71 Accessories for microtome
- ⁷2 Jars and stands for staining microscopy sections
- 73 Accessories for storing microscope sections
- 74 Brushes
- 75 Laboratory equipment for urine sediment analysis
- 75 Other laboratory equipment

Legend

- DG cargo in transport
- O DG cargo only in air transport
- temperature controlled cargo (transport regime 2-8°C)





Diagnostic and clinical products

The initial part of processing the clinical sample consists of high quality products necessary for further processing and final results of sample testing. That includes fixatives that contain the tissue's structure, alcohols, paraffins that enable section cutting and finally, immersion oils for microscopy and embedding media for long-term storage of stained sections.

- 8 Marking systems
- 10 Formalin fixatives
- 11 Other fixatives
- 13 Decalcifying solutions
- 14 Dehydrating and rehydrating agents
- 15 Clearing and deparaffinization agents
- 16 Mounting media
- 20 Immersion oils
- 21 Cleaning solutions for microscope lens and slides
- 22 Infiltration, embedding and demolding media

Marking systems

Marking systems consist of dyes for permanent tissue marking and markers used for marking glass slides, cassettes and transport containers. Tissue marking dyes are used for marking and orientation of tissue samples intended for further histopathology processing.

Tissue marking dyes, 5 color set

TMD <u>Tissue marking dyes</u> set 5x50 mL contains 5 dyes for tissue marking: green, yellow, black, red and blue. The set includes wooden sticks and rack.

5x50 mL 3408-SET

Tissue marking dyes, 7 color set

TMD <u>Tissue marking dyes</u> set 7x50 mL contains 7 dyes for tissue marking: green, yellow, black, red, blue, orange and lilac. The set includes wooden sticks and rack.

7x50 mL 3408-SET-7

Tissue marking dyes

For marking and orientation of tissue samples.

Tissue marking dyes, green	50 mL 100 mL 250 mL	3408-1-50 3408-1-100 3408-1-250
Tissue marking dyes, yellow	50 mL 100 mL 250 mL	3408-2-50 3408-2-100 3408-2-250
Tissue marking dyes, black	50 mL 100 mL 250 mL	3408-3-50 3408-3-100 3408-3-250
Tissue marking dyes, red	50 mL 100 mL 250 mL	3408-4-50 3408-4-100 3408-4-250
Tissue marking dyes, blue	50 mL 100 mL 250 mL	3408-5-50 3408-5-100 3408-5-250
Tissue marking dyes, orange	50 mL 100 mL 250 mL	3408-6-50 3408-6-100 3408-6-250
Tissue marking dyes, lilac	50 mL 100 mL 250 ml	3408-7-50 3408-7-100 3408-7-250



Duoline marker penProfessional marker pen for permanent labelling of cassettes, slides and transport containers. With fine and thick marker pen tip.

MD-1 1 pc

Liquid Blocker (PAP pen)Liquid Blocker contains hydrophobic liquid as a water repellent barrier around specimens or sections. Useful in immunohistochemistry that uses small amount of antiserum. Multiple specimens can be separated by drawn circles or lines applied to the same slide.

1 pc ML-1



Formalin fixatives

Fixation is one of the most important steps in tissue processing, preventing tissue degradation and decomposition of tissue and preserving properties important for detection of morphological and pathological changes in tissue samples.

4% NB Formaldehyde

10% neutral buffered and stabilized formalin solution, pH 7.0.

 1000 mL
 FNB4-1L

 5000 mL
 FNB4-5L

 10000 mL
 FNB4-10L

 20000 mL
 FNB4-20L

* 20 L volumes available in cubitainers on request

0

4% Formaldehyde

Stabilized 10% formalin solution.

 1000 mL
 F4-1L

 5000 mL
 F4-5L

 10000 mL
 F4-10L

 20000 mL
 F4-20L

0



10% NB Formaldehyde

25% neutral buffered and stabilized formalin solution, pH 7.0.

 1000 mL
 FNB10-1L

 5000 mL
 FNB10-5L

 10000 mL
 FNB10-10L

 20000 mL
 FNB10-20L

0

ImmunoForm

Stabilized 10% formalin solution produced from paraformaldehyde without methyl alcohol. For use ih histology and immunohistochemistry.

 1000 mL
 IMF-1L

 5000 mL
 IMF-5L

 10000 mL
 IMF-10L

0

10% Formaldehyde

Stabilized 25% formalin solution.

 1000 mL
 F10-1L

 5000 mL
 F10-5L

 10000 mL
 F10-10L

 20000 mL
 F10-20L

0

Formaldehyde 37% for use in histology

100% formalin solution stable at low temperatures.

 1000 mL
 F37H-1L

 5000 mL
 F37H-5L

 10000 mL
 F37H-10L

NB Formaldehyde concentrated 5x

Concentrated neutral buffered formalin for preparation of 10% buffered formalin (synonym: 4% buffered formaldehyde) solution (pH 7.0).

For preparation of 5 L of 4% buffered formaldehyde	1000 mL	FK5X-1L
For preparation of 25 L of 4% buffered formaldehyde	5000 mL	FK5X-5L
For preparation of 50 L of 4% buffered formaldehyde	10000 mL	FK5X-10L

Other fixatives

BioFix

Methyl alcohol-based fixative for cryostat sections.

500 mL BF-500 1000 mL BF-1L



BioFix C

Modified Carnoy fixative based on ethyl alcohol, chloroform and acetic acid.

 500 mL
 BFC-500

 1000 mL
 BFC-1L

 5000 mL
 BFC-5L



BioFix GL

Glyoxal-based fixative for use in histology.

500 mL BFGL-500 1000 mL BFGL-1L 5000 mL BFGL-5L

BioFix HD

Modified Hartmann/Davidson fixative that enables rapid fixation and better visualization of lymph nodes. The results provide clearer details of nucleus and minimize formalin pigment.

500 mL BFHD-500 1000 mL BFHD-1L 5000 mL BFHD-5L



BioForm S

Modified Schaeffer buffered fixative solution of formaldehyde in methanol.

1000 mL BIOS-OT-1L 5000 mL BIOS-OT-5L

Other fixatives

Bouin's solution

Aqueous solution of picric acid, formaldehyde and acetic acid for tissue fixation.

 100 mL
 BOU-OT-100

 1000 mL
 BOU-OT-1L

 5000 mL
 BOU-OT-5L

 10000 mL
 BOU-OT-10L

0

CitoFix

Solution for fixation and transport of cytological samples.

1000 mL CX-1L

CitoSpray

Solution for fixation of cytological smears in a spray bottle.

100 mL CS-100

CitoSpray solution

Solution for fixation of cytological smears in a bottle, for refill of CitoSpray bottle.

1000 mL CS-1L

DeHematin

Modified Kardasewitsch solution for depigmentation of tissues fixed in non-buffered formalin (after acid hematin occurs).

1000 mL KO-OT-1L 5000 mL KO-OT-5L

Saccomanno fixative

Solution for fixation and transport of cytology samples.

500 mL SF-OT-500 1000 mL SF-OT-1L 5000 mL SF-OT-5L



Decalcifying solutions

Decalcifiers most often contain inorganic or organic acids enabling rapid decalcification (i.e. calcium removal). Milder types of decalcifiers are formulated from chelating agents that are used for decalcification of more sensitive types of tissue.

OsteoFast 1

Light blue fixative and decalcifying solution for bone and hard tissue in histology, based on formaldehyde and hydrochloric acid.

1000 mL OF1-OT-1L 2500 mL OF1-OT-2.5L

OsteoFast 2

Light yellow decalcifying solution for bone and hard tissue in histology, based on formic acid.

1000 mL OF2-OT-1L 2500 mL OF2-OT-2.5L

OsteoSens

Decalcifying solution for sensitive calcified tissues in histology, contains EDTA.

1000 mL OS-OT-1L OS-OT-2.5L





Dehydrating and rehydrating agents

Alcoholic solutions and absolute alcohol are ideal for histocytology and cytology processing, storing and transport of samples. They are also used for dehydration and rehydration of tested samples.

Histanol 50

Denaturated 50% ethyl alcohol solution.

1000 mL H50-1L 5000 mL H50-5L 10000 mL H50-10L

Histanol 70

Denaturated 70% ethyl alcohol solution.

1000 mL H70-1L 5000 mL H70-5L 10000 mL H70-10L

Histanol 80

Denaturated 80% ethyl alcohol solution.

1000 mL H80-1L 5000 mL H80-5L 10000 mL H80-10L

Histanol 95

Denaturated 95% ethyl alcohol solution.

1000 mL H95-1L 5000 mL H95-5L 10000 mL H95-10L

Histanol 96

Denaturated 96% ethyl alcohol solution.

 1000 mL
 H96-1L

 5000 mL
 H96-5L

 10000 mL
 H96-10L

Histanol 100

Denaturated 100% ethyl alcohol.

 1000 mL
 H100-1L

 5000 mL
 H100-5L

 10000 mL
 H100-10L

Histanol G

Glycerol for use in microscopy.

1000 mL HG-1L 5000 mL HG-5L 10000 mL HG-10L

Histanol EP 95

Denaturated 95% mixture of ethyl and isopropyl alcohol solution.

1000 mL HEP95-1L 5000 mL HEP95-5L 10000 mL HEP95-10L

Histanol EP 100

Denaturated 100% mixture of ethyl and isopropyl alcohol.

 1000 mL
 HEP100-1L

 5000 mL
 HEP100-5L

 10000 mL
 HEP100-10L

Histanol IP

Absoute isopropyl alcohol for use in microscopy.

1000 mL HIP-1L 5000 mL HIP-5L 10000 mL HIP-10L

Histanol M

Absoute methyl alcohol for use in histology and cytology.

1000 mL HM-1L 5000 mL HM-5L 10000 mL HM-10L



Clearing and deparaffinization agents

Clearing agents are completely miscible with alcohol and paraffin, making them an ideal intermedium in histology and cytology processing. They are used multiple times during tissue processing and section staining; before embedding the tissue in paraffin, for removing the paraffin from the section before staining and before covering the stained section using medium for permanent preservation.

BioClear

Xylene-based medium for use in histology and cytology.

 1000 mL
 BC-1L

 2500 mL
 BC-2.5L

 5000 mL
 BC-5L

 10000 mL
 BC-10L

BioClear New

Xylene substitute. Based on aliphatic hydrocarbons, essentially no odor. For use in histology and cytology.

 1000 mL
 BCN-1L

 2500 mL
 BCN-2.5L

 5000 mL
 BCN-5L

 10000 mL
 BCN-10L



Mounting media

Ten different media for permanent covering of samples and mounting microscopy cover slips

In order to prepare permanent sections, stained samples must be covered with a suitable mounting medium. One of the most important criteria is the refractive index that must be similar to that of glass. Other criteria include a suitable solvent, polymer and medium viscosity. After the sample is covered with the medium and cover slide is mounted, the section is protected from external factors and remains preserved for a prolonged period of time. BioMount media do not crack nor turn yellow during sample storage.

BioGnost's synthetic BioMount media properties

NAME OF BIOMOUNT MEDIUM	SOLVENT	POLYMER	VISCOSITY @ 20°C (cSt, mPa)	REFRACTIVE INDEX @ 20°C (n _D)	USAGE IN AUTOMATED COVER SLIP MACHINES
BIOMOUNT DPX	xylene (BioClear)	S	850-950	1.5210-1.5230	-
BIOMOUNT DPX HIGH	xylene (BioClear)	S	1900-2100	1.5235-1.5255	-
BIOMOUNT DPX LOW	xylene (BioClear)	S	140-160	1.5115-1.5135	√
BIOMOUNT DPX NEW*	xylene (BioClear)	S	200-300	1.5150-1.5180	✓
BIOMOUNT NEW	isoparaffin (BioClear New)	М	950-1050	1.4385-1.4395	-
BIOMOUNT NEW LOW	isoparaffin (BioClear New)	М	130-160	1.4345-1.4365	√
BIOMOUNT	xylene (BioClear)	М	450-650	1.4920-1.4930	✓
BIOMOUNT HIGH	xylene (BioClear)	М	5000-5500	1.4900-1.4920	-
BIOMOUNT M	toluene	М	140-160	1.4900-1.4920	√
BIOMOUNT C	xylene (BioClear)	М	5000-5500	1.4910-1.4925	-

Legend: M - methacrylate, S - styrene (polystyrene)

Water-based mounting media

BIOMOUNT AQUA	water	PVP	1800-2100	1.3630-1.3670	-
NAME OF BIOMOUNT MEDIUM	SOLVENT	POLYMER	VISCOSITY @ 20°C (cSt, mPa)	REFRACTIVE INDEX @ 20°C (n _D)	AUTOMATED COVER SLIP MACHINES

LICACE IN

 $^{^{\}star}$ does not contain harmful dibutyl phthalate plasticizer

 $[\]ensuremath{^{**}}$ contact us to find out how to decrease viscosity of a medium

Xylene and styrene-based media for permanent mounting

BioMount DPX

Medium viscosity medium for permanent covering of microscope samples and mounting cover slips.

100 mL BM-100 250 mL BM-250 500 mL BM-500



BioMount DPX High

High viscosity medium for permanent covering of microscope samples and mounting cover slips.

 100 mL
 BMH-100

 250 mL
 BMH-250

 500 mL
 BMH-500



BioMount DPX Low

Low viscosity medium for permanent covering of microscope samples and mounting cover slips. Can be used in automatic cover slip machines.

 100 mL
 BML-100

 250 mL
 BML-250

 500 mL
 BML-500





BioMount DPX New

Low viscosity medium for permanent covering of microscope samples and mounting cover slips. Medium contains more environmentally friendly plasticizer instead of harmful dibutyl phthalate. Can be used in automatic cover slip machines.

 100 mL
 BMLE-100

 250 mL
 BMLE-250

 500 mL
 BMLE-500

Methacrylate and xylene substitute-based media for permanent mounting

BioMount New

Medium viscosity for permanent covering of microscope samples deparaffinized using xylene substitute-based medium, as well as xylene-based medium.

30 mL BMN-30 100 mL BMN-100 250 mL BMN-250 500 mL BMN-500

BioMount New Low

Very low viscosity for permanent covering of microscope samples deparaffinized using xylene substitute-based medium, as well as xylene-based medium.

30 mL	BMNL-30
100 mL	BMNL-100
250 mL	BMNL-250
500 mL	BMNL-500



Mounting media

Methacrylate-based permanent mounting media

BioMount

Low viscosity xylene-based medium for permanent covering of microscope slides and mounting cover slips.

 100 mL
 BMT-100

 250 mL
 BMT-250

 500 mL
 BMT-500



BioMount High

Very high viscosity xylene-based medium for permanent covering of microscope samples and mounting cover slips.

 100 mL
 BMTH-100

 250 mL
 BMTH-250

 500 mL
 BMTH-500

BioMount M

Very low viscosity toluene-based medium for permanent covering of microscope samples and mounting cover slips.

100 mL BMM-100 250 mL BMM-250 500 mL BMM-500





Additional permanent mounting media

Canada Balsam

Very high viscosity natural medium for permanent covering of microscope slides and mounting cover slips.

 100 mL
 KB-100

 250 mL
 KB-250

 500 mL
 KB-500

* in order to conserve and protect our environment, we advise using **BioMount C** synthetic medium

BioMount C

Very high viscosity medium for permanent covering of microscope slides and mounting cover slips.

100 mL BMC-100 250 mL BMC-250 500 mL BMC-500

* excellent synthetic substitution for natural covering media, such as Canada Balsam





Water-based mounting media

BioMount Aqua

Water-based medium for covering microscope slides and mounting cover slips. With application in immunohistochemistry as a barrier between chromogen stains and the organic solvent-based permanent mounting media.

30 mL	BMA-30
100 mL	BMA-100
250 mL	BMA-250
500 mL	BMA-500

Immersion oils

Seven different types of immersion oils for microscopical analysis

Immersion oils are essential for routine laboratory work, especially in hematology, cytology and microbiology. They are commonly used for high scale magnifications in order to secure high resolution and a clear image. There are several types of immersion oils that differ according to viscosity and certain properties required by different microscopy techniques. They all have a common refractive index of 1.5150, equal to the refractive index of glass.

BioGnost's Immersion oils properties

NAME OF	VISCOSITY	COMBINING WITH	TYPE OF MICROSCOPY
IMMERSION OIL	@ 23°C (cSt, mPa)	OTHER TYPES OF OIL	TYPE OF MICROSCOPY
IMMERSION OIL	800-1300	Immersion oil type A Immersion oil type C	light
IMMERSION OIL TYPE A	100-300	Immersion oil Immersion oil type C	light
CEDARWOOD OIL	1500-2200	-	light
IMMERSION OIL TYPE C	1500-2200	Immersion oil Immersion oil type A	light
IMMERSION OIL TYPE FF	170	-	fluorescence
IMMERSION OIL TYPE 37*	1250**	-	light, for use at 37°C
IMMERSION OIL, TROPICAL GRADE***	1000-1500****	-	light, for use at 27-29°C
* refractive index n _p at 37°C = 1.5150	** viscosity at 37°C *** re	fractive index n_n at tropical temperatures =	1.515 **** viscosity at 28°C

Immersion oil

Immersion oil for microscopical analysis with optimal viscosity of 600-1100 cSt.

10x10 mL	IU-1010
30 mL	IU-30
100 mL	IU-100
250 mL	IU-250
500 mL	IU-500



Immersion oil for microscopical analysis with low viscosity of 100-300 cSt.

10x10 mL	IUA-1010
30 mL	IUA-30
100 mL	IUA-100
250 mL	IUA-250
500 mL	IUA-500





Cedarwood oil

Natural immersion oil for microscopical analysis with high viscosity of 1500-2200 cSt.

100 mL CU-100 250 mL CU-250 500 mL CU-500

* in order to conserve and protect our environment, we advise using **Immersion oil type C**

Immersion oil type C

Immersion oil for microscopical analysis with high viscosity of 1500-2200 cSt.

100 mL IUC-100 250 mL IUC-250 500 mL IUC-500

* excellent synthetic substitution for natural immersion oils, such as **Cedarwood oil**

Immersion oil type FF

Immersion oil for microscopical analysis in fluorescence microscopy because of complete absence of fluorescence.

100 mL IUF-100 250 mL IUF-250 500 mL IUF-500

Immersion oil type 37

Immersion oil for microscopical analysis with optimal refractive index and viscosity of 1250 cSt at 37°C.

 100 mL
 IU37-100

 250 mL
 IU37-250

 500 mL
 IU37-500



Immersion oil, tropical grade

Immersion oil for microscopical analysis in tropical climates with optimal refractive index at 27-29°C.

 10x10 mL
 IUT-1010

 30 mL
 IUT-30

 100 mL
 IUT-100

 250 mL
 IUT-250

 500 mL
 IUT-500





Cleaning solutions for microscope lens and slides

Clean Lens, solution 1

Solution according to Carl Zeiss formulation for cleaning microscope lens. Based on n-hexane and 2-propanol.

100 mL CL1-OT-100



Clean Lens, solution 2

Solution for cleaning microscope lens and removing immersion oil from stained slide after using microscope. Based on diethyl ether and 2-propanol.

100 mL CL2-OT-100

Clean Lens, solution 3

Solution for cleaning microscope lens and removing immersion oil from stained slide after using microscope. Based on petroleum ether and 2-propanol.

100 mL CL3-OT-100

Infiltration, embedding and demolding media

Paraffin is the most commonly used medium for infiltration, embedding and forming of tissue blocks. It provides support to the tissue block during cutting of thin histology sections.

BioWax 52/54

Blend of paraffin waxes with low melting point at 52/54°C for processing wide range of different samples. Lower melting point makes it suitable for working with delicate tissues without deforming or damaging and allows good preservation of tissue morphology during processing.

1 kg	BW52/54-1
2 kg	BW52/54-2
10 kg	BW52/54-10

BioWax 56/58

Blend of paraffin waxes with melting point at 56/58°C for optimal infiltration and routine work, enables easier formation of ribbons, virtually no cut resistance and better overall results.

1 kg	BW56/58-1
2 kg	BW56/58-2
10 kg	BW56/58-10

BioWax Blue

Blend of blue paraffin waxes with melting point at 56/58°C and added plastic polymers for optimal infiltration. Especially suitable for small tissue samples. Blue color provides the contrast that enables easier visualization during processing. It is also recommended for use in immunoperoxidase methods.

1 kg	BWB-1
2 kg	BWB-2
10 kg	BWB-10

BioWax Micro

Blend of paraffin waxes with melting point at 56/58°C for optimal infiltration and routine work. Microcrystalline additive allows high quality tissue infiltration, easy separation of the block, flexibility and excellent section shaping without deformation.

1 kg	BWM-1
2 kg	BWM-2
10 ka	BWM-10

BioWax Plus 56/58

An optimal blend of paraffin wax and plastic polymers with melting point at 56/58°C. Its properties include low shrinking rate during cooling down and excellent section creation without deformations. Provides excellent infiltration into tissues and reduces the tendency of tissue to crack.

1 kg	BWPLUS-1
2 kg	BWPLUS-2
10 kg	BWPLUS-10

Micro Clean

Solution in spray bottle for rapid and efficient paraffin residue removal from work surfaces and instruments.

100 mL MC-OT-100





CryoFix Gel

Embedding matrix for cryosectioning of tissue samples known as OCT medium (optimal cutting temperature). Efficient at temperatures of -10°C and below. Available in 6 different colors.

CryoFix Gel	100 mL	CF-100
CryoFix Gel pink	100 mL	CFR-100
CryoFix Gel blue	100 mL	CFP-100
CryoFix Gel green	100 mL	CFZ-100
CryoFix Gel yellow	100 mL	CFY-100



Cryo Spray Medium in spray bottle for freezing histological samples.

200 mL 08-SPRAY-2

DeMold Spray

Solution in spray bottle for easy removal of paraffin tissue blocks from the mold.

100 mL DS-100

DeMold solution

Solution for refilling DeMold Spray bottle.


500 mL DS-500 1000 mL DS-1L

Tissue Soft

Solution for softening the tissue embedded in paraffin block before microtome sectioning.

500 mL TS-500 1000 mL TS-1L







Standard and special diagnostic stains

Diagnostic reagents for staining, demonstrating and contrasting of samples, as well as for achieving superb staining results in all fields of clinical pathology (histology, cytology, hematology, bacteriology). It is possible to order certified powder dyes for individual preparation of reagents.

- 26 Histology stains
- 30 Cytology stains
- 32 Hematology stains
- 36 Bacteriology stains
- 41 Fungal stains
- 42 Special stains for histology and cytology
- 58 Powder dyes and stains for use in microscopy


Hematoxylin reagents

Nuclear stains for standard HE staining

Properties of BioGnost's Hematoxylin reagents

TYPE OF HEMATOXYLIN	CYTOLOGY	HISTOLOGY	IHC	PROGRESSIVE METHOD	REGRESSIVE METHOD
HEMATOXYLIN H	-	✓	-	✓	√
HEMATOXYLIN M	-	✓	✓	✓	-
HEMATOXYLIN ML	-	✓	✓	✓	√
HEMATOXYLIN G1	√	✓	✓	✓	-
HEMATOXYLIN G2	√	✓	✓	✓	-
HEMATOXYLIN G3	✓	✓	-	✓	-
HEMATOXYLIN CAR	-	✓	✓	✓	-
HEMATOXYLIN COL	-	✓	✓	✓	√
HEMATOXYLIN HP, PAP 1A	✓	-	-	✓	√
HEMATOXYLIN HP, PAP 1B	✓	-	-	✓	√
HEMATOXYLIN W KIT	-	✓	✓	✓	acid resistant
HEMATOXYLIN P.T.A.	-	✓	-	✓	-

Each of BioGnost's Hematoxylins is specially stabilized in order to maintain the reagent's reactivity for prolonged periods of time, as well as to maintain the quality of the reagent. Every staining provides a crystal clear and highly detailed image of the cell's nucleus, which in turn makes abnormal or pathological changes easier to detect.





Hematoxylin H

Modified hematoxylin according to Harris, most commonly used formulation in HE staining. Strong intensity reagent for progressive and regressive staining in histopathology.

100 mL	HEMH-OT-100
500 mL	HEMH-OT-500
1000 mL	HEMH-OT-1L
2500 mL	HEMH-OT-2.5L

Hematoxylin M

Modified hematoxylin according to Mayer. Weaker intensity reagent for progressive staining in histopathology and counterstaining in immunohistochemistry.

100 mL	HEMM-OT-100
500 mL	HEMM-OT-500
1000 mL	HEMM-OT-1L
2500 mL	HEMM-OT-2.5L

Hematoxylin ML

Modified hematoxylin according to Mayer-Lillie. Strong intensity new generation reagent for progressive and regressive staining in histopathology.

100 mL	HEMML-OT-100
500 mL	HEMML-OT-500
1000 mL	HEMML-OT-1L
2500 mL	HEMML-OT-2.5L

Hematoxylin G1

Hematoxylin according to Gill, ideal for staining goblet cells. New generation reagent for delicate progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG1-OT-100
500 mL	HEMG1-OT-500
1000 mL	HEMG1-OT-1L
2500 mL	HEMG1-OT-2.5L

Hematoxylin G2

Hematoxylin according to Gill, contains double concentration of hematoxylin compared to Hematoxylin G1. Moderate intensity new generation reagent for progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG2-OT-100
500 mL	HEMG2-OT-500
1000 mL	HEMG2-OT-1L
2500 mL	HEMG2-OT-2.5L

Hematoxylin G3

Modified hematoxylin contains triple concentration of hematoxylin compared to Hematoxylin G1. Strong intensity new generation reagent for progressive staining in histopathology and cytology.

100 mL	HEMG3-OT-100
500 mL	HEMG3-OT-500
1000 mL	HEMG3-OT-1L
2500 mL	HEMG3-OT-2.5L



Hematoxylin reagents

Nuclear stains for standard HE staining

Hematoxylin W kit

Acid-resistant hematoxylin according to Weigert. Two-reagent kit that stains the nuclei blue-black, often a component of special staining kits for connective tissues.

2x100 mL HEMW-K-100 2x500 mL HEMW-K-500 2x1000 mL HEMW-K-1L



Hematoxylin P.T.A.

Synonyms: Hematoxylin Phosphotungstic Acid, Mallory P.T.A.H. reagent. Simultaneously stains different types of tissue reddishbrown and blue. For differentiation between fibrin, smooth and cross-striated muscle fibers.

500 mL HPTA-OT-500 1000 mL HPTA-OT-1L

Hematoxylin CAR

Modified hematoxylin according to Carazzi for delicate nuclear staining without staining cytoplasmic components. Reagent for progressive staining of frozen sections in histopathology.

 100 mL
 HEMCA-OT-100

 500 mL
 HEMCA-OT-500

 1000 mL
 HEMCA-OT-1L

Hematoxylin COL

Modified hematoxylin according to Cole, contains iodine. Weaker intensity reagent for progressive and regressive staining in histopathology.

 100 mL
 HEMCO-OT-100

 500 mL
 HEMCO-OT-500

 1000 mL
 HEMCO-OT-1L

Hematoxylin Instant kit

Two-component kit that contains ready-to-use powders for preparation of hematoxylin reagent for progressive and regressive nuclear staining in histopathology and cytology.

for 1000 mL HEMI-K-1L for 6000 mL HEMI-K-6L

Kit for rapid HE staining of frozen and paraffin-embedded sections

Ready-to-use eight-reagent kit (in 20 containers that can be used as staining jars) for rapid HE staining of frozen and paraffin tissue sections in histopathology. Contains xylene substitute as clearing agent and xylene substitute-based medium for permanent coverslipping.

for 100 tests HE-RTU-100T



Differentiation and bluing reagents for histology and cytology

Differentiation of the nucleus enables clear visualization of nuclear structures in regressive procedure, while the bluing reagent turns the nucleus blue.

Acid alcohol

Reagent for enabling crisp visualization of nuclear chromatin detail and removing of excess background dye in histological samples after regressive staining with hematoxylin.

1000 mL KA-OT-1L 2500 mL KA-OT-2.5L



HCL Pap reagent

Buffered reagent for enabling crisp visualization of nuclear chromatin detail and removing of excess background dye in cytological after regressive staining with hematoxylin.

1000 mL HCLPAP-OT-1L 2500 mL HCLPAP-OT-2.5L

Scott's solution

Buffered reagent for conversion of red hematoxylin stained nuclei to optimal blue.

1000 mL SC-OT-1L 2500 mL SC-OT-2.5L

Bluing reagent

Buffered reagent for conversion of red hematoxylin stained nuclei to optimal blue.

1000 mL BR-OT-1L 2500 mL BR-OT-2.5L

Eosin solutions

Cytoplasmic stains for standard HE staining

Eosin Y (yellowish eosin) solutions are commonly used as a counterstain to hematoxylin in the standard histology HE staining method. Eosin Y is an anionic dye that stains basic cell components (such as cytoplasm, collagen, muscle fibers and erythrocytes) bright red.

Eosin Y aqueous solutions

Eosin Y aqueous solutions for cytoplasmic counterstaining.

Eosin Y 0.2% aqueous	1000 mL 2500 mL	EOY-02-OT-1L EOY-02-OT-2.5L
Eosin Y 0.5% aqueous	1000 mL 2500 mL	EOY-05-OT-1L EOY-05-OT-2.5L
Eosin Y 1% aqueous	1000 mL 2500 mL	EOY-10-OT-1L EOY-10-OT-2.5L
Eosin Y 2% aqueous	1000 mL 2500 mL	EOY-20-OT-1L EOY-20-OT-2.5L

Alcoholic Eosin Y solutions

Alcoholic eosin Y solutions for cytoplasmic counterstaining.

Eosin Y 0.5% alcoholic	1000 mL 2500 mL	EOYA-05-OT-1L EOYA-05-OT-2.5L
Eosin Y 1% alcoholic	1000 mL 2500 mL	EOYA-10-OT-1L EOYA-10-OT-2.5L
Eosin Y 2% alcoholic	1000 mL 2500 mL	EOYA-20-OT-1L EOYA-20-OT-2.5L

Eosin Contrast

Modified alcoholic solution for cytoplasmic counterstaining. The reagent contains Eosin Y and Phloxine B for enhanced counterstaining effect.

	L
2500 mL EOYK-OT-2.	5L



Eosin Contrast PLUS

Modified alcoholic solution for intensive cytoplasmic counterstaining. Reagent contains eosin Y, Phloxine B, and Biebrich Scarlet dyes for additional counterstaining effect.

1000 mL	EOYKP-OT-1L
2500 mL	EOYKP-OT-2.5L



Eosin Y instant

Mixture of powders for preparation of eosin reagent in aqueous or alcoholic medium.

Eosin Y Instant, aqueous	for 1000 mL	EOYI-AQ-1L
Eosin Y Instant,	for 1000 mL	EOYI-ALC-1L





Papanicolaou (Pap) reagents

Nuclear and cytoplasmic reagents for Pap staining

Because of their simplicity of use and exceptional precision in diagnostics, reagents for staining according to Papanicolaou have substantially contributed to decreasing occurrence of cervical cancer. By using a simple staining protocol, BioGnost's Pap reagents enable excellent differentiation between cellular structures.

Hematoxylin HP, Pap 1A

Modified Harris hematoxylin solution for nuclear staining according to Papanicolaou. Strong intensity reagent for progressive and regressive staining in exfoliative cytology.

100 mL	HEMHP-OT-100
500 mL	HEMHP-OT-500
1000 mL	HEMHP-OT-1L
2500 mL	HEMHP-OT-2.5L

Hematoxylin HP, Pap 1B

Hematoxylin modified according to Harris for intensive nuclear staining according to Papanicolaou. Reagent for progressive and regressive staining in exfoliative cytology, with formulation stronger than Hematoxylin HP, Pap 1A.

100 mL	HEMHP1B-OT-100
500 mL	HEMHP1B -OT-500
1000 mL	HEMHP1B -OT-1L
2500 ml	HEMHP1R -OT-2 5I

Kit for rapid Pap staining

Ready-to-use eight-reagent kit (in 20 containers that can be used as staining jars) for rapid progressive gynecology and non-gynecology cytological samples. Contains xylene substitute as clearing agent and xylene substitute-based medium for permanent covering of samples.

for 100 tests PAP-100T




Shorr's solution

Enables clear differentiation of eosinophilic and cyanophilic cells of cervical epithelium. It is also used for a better visualization of sperm morphology.

100 mL	SH-OT-100
500 mL	SH-OT-500
1000 mL	SH-OT-1L
2500 mL	SH-OT-2.5L





Hematoxylin G1

Hematoxylin according to Gill, ideal for staining goblet cells. New generation reagent for delicate progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG1-OT-100
500 mL	HEMG1-OT-500
1000 mL	HEMG1-OT-1L
2500 mL	HEMG1-OT-2.5L

Hematoxylin G2

Hematoxylin according to Gill, contains double concentration of hematoxylin compared to Hematoxylin G1. Moderate intensity new generation reagent for progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG2-OT-100
500 mL	HEMG2-OT-500
1000 mL	HEMG2-OT-1L
2500 ml	HFMG2-OT-2 5I

Hematoxylin G3

Modified hematoxylin contains triple concentration of hematoxylin compared to Hematoxylin G1. Strong intensity new generation reagent for progressive staining in histopathology and cytology.

100 mL	HEMG3-OT-100
500 mL	HEMG3-OT-500
1000 mL	HEMG3-OT-1L
2500 mL	HEMG3-OT-2.5L

OG-6 reagent, Pap 2A

Monochromatic reagent for counterstaining mature and keratinized squamous cell. Contains Orange G dye that stains the cytoplasm orange.

100 mL	OG6-OT-100
500 mL	OG6-OT-500
1000 mL	OG6-OT-1L
2500 mL	OG6-OT-2.5L



Orange II reagent, Pap 2B

Monochromatic reagent for counterstaining mature and keratinized squamous cell. Contains Orange II dye that stains the cytoplasm intensive red.

100 mL	OR2-OT-100
500 mL	OR2-OT-500
1000 mL	OR2-OT-1L
2500 mL	OR2-OT-2.5L



OG-EA31 Pap reagent

One-step polychromatic reagent used as counterstain for gynecology samples in cytology. Enables rapid staining and differentiation between epithelial cells whose cytoplasm is stained green and pink. It is not necessary to use OG-6, Pap 2A reagent during staining because it stains cytoplasm of keratinized cell orange at the same time.

 100 mL
 OGEA31-OT-100

 500 mL
 OGEA31-OT-500

 1000 mL
 OGEA31-OT-1L

 2500 mL
 OGEA31-OT-2.5L

OG-EA50 Pap reagent

One-step polychromatic reagent used as counterstain for gynecology samples in cytology. Enables rapid staining and differentiation between epithelial cells whose cytoplasm is stained blue and pink. It is not necessary to use OG-6, Pap 2A reagent during staining because it stains cytoplasm of keratinized cell orange at the same time.

 100 mL
 OGEA50-OT-100

 500 mL
 OGEA50-OT-500

 1000 mL
 OGEA50-OT-1L

 2500 mL
 OGEA50-OT-2.5L



EA 31 reagent, Pap 3A

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **gynecology** samples in cytology, contains the largest amount of green dye compared to other EA reagents. Stains the cytoplasm blue-green and pink, with emphasis on green and pink color.

 100 mL
 EA31-OT-100

 500 mL
 EA31-OT-500

 1000 mL
 EA31-OT-1L

 2500 mL
 EA31-OT-2.5L



Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **gynecology** samples in cytology, stains the cytoplasm blue-green and pink, with emphasis on blue and pink color.

 100 mL
 EA50-OT-100

 500 mL
 EA50-OT-500

 1000 mL
 EA50-OT-1L

 2500 mL
 EA50-OT-2.5L



EA 65 reagent, Pap 3C

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **non-gynecology** samples in cytology, stains the cytoplasm red.

 100 mL
 EA65C-OT-100

 500 mL
 EA65C-OT-500

 1000 mL
 EA65C-OT-1L

 2500 mL
 EA65C-OT-2.5L



EA 65 reagent, Pap 3D

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **non-gynecology** samples in cytology, stains the cytoplasm light blue-green and pink.

 100 mL
 EA65D-OT-100

 500 mL
 EA65D-OT-500

 1000 mL
 EA65D-OT-1L

 2500 mL
 EA65D-OT-2.5L





Bio-Diff kits

Rapid staining solutions for hematology, cytology and histology

Rapid diagnostic kits that provide precise staining results in a short period of time, such as results provided by the May-Gruenwald Giemsa (MGG) staining method. Each component of Bio-Diff and Bio-Diff RTU kits is stabilized and prepared according to the highest standards.

Bio-Diff kit

Three-reagent kit that contains fixative agent, red and blue components for fast and effective staining. Each kit contains buffer tablets for consistent staining results.

3x100 mL	BD-K-100
3x500 mL	BD-K-500
3x1000 mL	BD-K-1L
3x2500 mL	BD-K-2.5L



Bio-Diff 1 reagent

 Fixative agent, component of Bio-Diff kit.

 100 mL
 BD1-OT-100

 500 mL
 BD1-OT-500

 1000 mL
 BD1-OT-1L

 2500 mL
 BD1-OT-2.5L

Bio-Diff 2 reagent

Buffered red component of Bio-Diff kit.

100 mL	BD2-OT-100
500 mL	BD2-OT-500
1000 mL	BD2-OT-1L
2500 mL	BD2-OT-2.5L

Bio-Diff 3 reagent

Buffered blue component of Bio-Diff kit.

100 mL	BD3-OT-100
500 mL	BD3-OT-500
1000 mL	BD3-OT-1L
2500 mL	BD3-OT-2.5L







Bio-Diff RTU kit

Ready-to-Use three-reagent kit with reagents stored in containers that can be used as staining jars. Kit contains fixative agent, red and blue components for fast and effective staining and buffer tablet for consistent staining results.

3x100 mL BD-RTU-100



Bio-Diff 1 RTU reagent

Fixative component Bio-Diff RTU kit.

100 mL BD1-RTU

Bio-Diff 2 RTU reagent

Buffered red component of Bio-Diff RTU kit.

100 mL BD2-RTU

Bio-Diff 3 RTU reagent

Buffered blue component of Bio-Diff RTU kit.

100 mL BD3-RTU



Buffer solution, pH 7.2

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

 1000 mL
 PUF72-OT-1L

 2500 mL
 PUF72-OT-2.5L

 5000 mL
 PUF72-OT-5L

Buffer tablets pH 7.2

Phosphate buffer tablets for preparing buffer solutions used in hematology and histology. One buffer tablet is sufficient for preparation of 1000 mL of phosphate buffer solution.

50 pcs PT-72-50 100 pcs PT-72-100

Use buffer pH 6.8 for hematological and cytological diagnostics. Use buffer pH 7.2 for the detection of blood parasites (see page 34).



May-Gruenwald and Giemsa reagents

Romanowsky staining solutions for hematology, cytology and histology

One of the most commonly used methods (Pappenheim) is May-Gruenwald Giemsa (MGG) for staining hematology and cytology materials, as well as cytology smears of hematopoietic organs. Giemsa solution is also used for staining histology sections.

Buffer solution, pH 6.4

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

 1000 mL
 PUF64-OT-1L

 2500 mL
 PUF64-OT-2.5L

 5000 mL
 PUF64-OT-5L

Buffer solution, pH 6.8

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

 1000 mL
 PUF68-OT-1L

 2500 mL
 PUF68-OT-2.5L

 5000 mL
 PUF68-OT-5L

Buffer tablets pH 6.8

Phosphate buffer tablets for preparing buffer solutions used in hematology and histology. One buffer tablet is sufficient for preparation of 1000 mL of phosphate buffer solution.

50 pcs PT-68-50 100 pcs PT-68-100

Use buffer pH 6.8 for hematological and cytological diagnostics. Use buffer pH 7.2 for the detection of blood parasites (see page 33).

May-Gruenwald solution

Polychromatic fixative solution containing Eosin Y, Methylene Blue and azure dyes. For staining in hematology, cytology and staining sections of hematopoietic organs in histopathology.

 100 mL
 MG-OT-100

 10x100 mL
 MG-OT-110

 500 mL
 MG-OT-500

 1000 mL
 MG-OT-1L

 2500 mL
 MG-OT-2.5L



Giemsa solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For use in hematology, cytology, histology and for staining section samples of hematopoietic organs.

 100 mL
 GM-OT-100

 10x100 mL
 GM-OT-110

 500 mL
 GM-OT-500

 1000 mL
 GM-OT-1L

 2500 mL
 GM-OT-2.5L





Additional staining solutions for hematology and cytology

Brilliant Cresyl Blue, solution

Solution for staining reticulocytes in blood.

 100 mL
 BCB-OT-100

 500 mL
 BCB-OT-500

 1000 mL
 BCB-OT-1L

Hayem's solution

Solution for manual counting of erythrocytes.

100 mL HY-OT-100 500 mL HY-OT-500 1000 mL HY-OT-1L

Tuerk's solution

Solution for manual counting of leukocytes.

100 mL TU-OT-100 500 mL TU-OT-500 1000 mL TU-OT-1L



Leishman's solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For staining in hematology and clinical cytology.

 100 mL
 LE-OT-100

 500 mL
 LE-OT-500

 1000 mL
 LE-OT-1L



Wright's solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For staining in hematology, cytology and cytogenetics.

 100 mL
 WR-OT-100

 500 mL
 WR-OT-500

 1000 mL
 WR-OT-1L





Field kit

Ready to use two-reagent kit for rapid and efficient staining and detection of parasites in hematology samples. Kit for 100 tests contains reagents stored in containers that can be used as staining jars.

for 100 tests FD-100T 2x500 mL FD-K-500

Wright-Giemsa solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. Modified Romanowsky stain for hematology staining.

 100 mL
 WRGM-OT-100

 500 mL
 WRGM-OT-500

 1000 mL
 WRGM-OT-1L

 2500 mL
 WRGM-OT-2.5L



Wright-Giemsa, solution for reptiles

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. Modified Romanowsky dye for staining in reptile hematology.

 100 mL
 WRGG-OT-100

 500 mL
 WRGG-OT-500

 1000 mL
 WRGG-OT-1L

 2500 mL
 WRGG-OT-2.5L



BioGram kits

Kits for identification of bacteria according to Gram

Differential staining of bacteria according to Gram is widely known and used method in microbiology. BioGram 4 kit uses standard method to provide optimal staining and identification of Gram-positive and Gram-negative bacteria. BioGram ECO kit is a phenol-free, environment-friendly alternative for BioGram 4 kit with equivalent quality result of staining.

BioGram 4 kit

Four-reagent kit for identification of bacteria according to Gram. Kit contains Gram Crystal Violet 1% solution, stabilized Gram Lugol solution, double amount of Gram Decolorizer solution 2 and Gram Safranin solution as counterstain.

for 100 tests	BGR4-100T
5x100 mL	BGR4-K-100
5x250 mL	BGR4-K-250
5x500 mL	BGR4-K-500



Gram Crystal Violet 1% solution

Primary stain reagent for Gram-positive bacteria. Crystal Violet is also known as Gentian Violet and Methyl Violet 10B.

GC1-OT-100
GC1-OT-250
GC1-OT-500
GC1-OT-1L
GC1-OT-2.5L



Gram Lugol solution, stabilized

Solution contains polyvinylpyrrolidone iodine (PVP-I).

100 !	CLC OT 100
100 mL	GLS-OT-100
250 mL	GLS-OT-250
500 mL	GLS-OT-500
1000 mL	GLS-OT-1L
2500 ml	GLS-OT-2 51

Gram Decolorizer solution 2

Medium speed decolorizer used after stabilized iodine solution.

100 mL	GD2-OT-100
250 mL	GD2-OT-250
500 mL	GD2-OT-500
1000 mL	GD2-OT-1L
2500 mL	GD2-OT-2.5L



Gram Safranin solution

Counterstain for Gram-negative bacteria, contains Safranin O dye.

100 mL 250 mL 500 mL 1000 mL	GSF-OT-100 GSF-OT-250 GSF-OT-500 GSF-OT-1L
2500 mL	GSF-OT-2.5L
2500 mL	GSF-OT-2.5L



BioGram ECO kit

Four-reagent **phenol-free** kit for the identification of bacteria according to Gram. Kit contains Gram Crystal violet, phenol free reagent, Gram Sodium hydrogencarbon, solution, stabilized Gram Lugol solution, double amount of Gram Decolorizer solution 2 and Gram Safranin solution as counterstain.

2x50 mL+4x100 mL	BGRE-K-100
2x250 mL+4x500 mL	BGRE-K-500



Gram Crystal Violet, phenol-free reagent

Primary stain reagent for Gram-positive bacteria, doesn't contain phenol. Crystal Violet is also known as Gentian Violet and Methyl Violet 10B.

100 mL	GCF-OT-100
250 mL	GCF-OT-250
500 mL	GCF-OT-500
1000 mL	GCF-OT-1L
2500 mL	GCF-OT-2.5L



Gram sodium hydrogencarbonate, solution

Solution of sodium hydrogencarbon for use with BioGram ECO kit.

100 mL	GNHK-OT-100
250 mL	GNHK-OT-250
500 mL	GNHK-OT-500
1000 mL	GNHK-OT-1L
2500 mL	GNHK-OT-2.5L

Additional bacteriology staining solutions according to Gram

Gram Lugol solution

Non-stabilized Lugol solution.

100 mL	GL-OT-100
250 mL	GL-OT-250
500 mL	GL-OT-500
1000 mL	GL-OT-1L

Gram Decolorizer solution 1

Slow acting decolorizer solution used after non-stabilized iodine solution.

100 mL	GD1-OT-100
250 mL	GD1-OT-250
500 mL	GD1-OT-500
1000 mL	GD1-OT-1L



Gram Decolorizer solution 3

Fast acting solution used with automated stainers.

GD3-OT-100
GD3-OT-250
GD3-OT-500
GD3-OT-1L

Gram Safranin 5x concentrated

Concentrated counterstain solution of Safranin O dye.

For preparation of 500 mL of Safranin O counterstain solution	100 mL	SO05X-OT-100
For preparation of 2500 mL of Safranin O counterstain solution	500 mL	SO05X-OT-500
For preparation of 5000 mL of Safranin O counterstain solution	1000 mL	SO05X-OT-1L



Carbol Fuchsin Gram 10x concentrated

Concentrated Carbol Fuchsin solution for preparation of counterstain solution for modified staining. Also known as concentrated Gram Fuchsin solution.

For preparation of 1000 mL of Carbol Fuchsin Gram counterstain solution	100 mL	CFGR10-OT-100
For preparation of 5000 mL of Carbol Fuchsin Gram counterstain solution	500 mL	CFGR10-OT-500
For preparation of 10000 mL of Carbol Fuchsin Gram counterstain solution	1000 mL	CFGR10-OT-1L



Additional iodine solutions

Lugol's solution (Schiller's test)

Classic formulation of aqueous solution of iodine and potassium iodide. Commonly used in gynecology.

100 mL LUG-OT-100 250 mL LUG-OT-250 500 mL LUG-OT-500

Lugol's C solution

Aqueous solution of iodine and potassium iodide, enhanced formulation with stronger staining (compared to classic Lugol solution).

 100 mL
 LUGC-OT-100

 250 mL
 LUGC-OT-250

 500 mL
 LUGC-OT-500

Lugol's solution, acid

Aqueous iodine and potassium iodide solution with added acid.

100 mL	LUGK-OT-100
250 mL	LUGK-OT-250
500 mL	LUGK-OT-500

TB-Stain Hot and TB-Stain ECO kits

Kits for staining acid-fast bacteria according to Ziehl-Neelsen

Most commonly known method for detecting TB bacteria according to Ziehl-Neelsen. This method enables rapid and easy staining of acid fast bacterial strains, such as *Mycobacteria*, *Nocardia*, *Rhodococcus*, *Dietzia* and *Gordonia*. TB-Stain ECO kit is a phenol–free, environment-friendly alternative for TB-Stain Hot kit.

TB-Stain Hot kit

Three-reagent kit for staining acid-fast bacteria. Contains TB Carbol Fuchsin reagent, double amount of TB Decolorizer and Methylene Blue Loeffler's reagent as counterstain.

for 100 tests	TBH-100T
4x100 mL	TBH-K-100
4x250 mL	TBH-K-250
4x500 mL	TBH-K-500



TB Carbol Fuchsin reagent

Synonym: Ziehl-Neelsen reagent. Primary stain reagent for acid-fast bacteria used in TB-Stain Hot, TB-Stain Cold and TB-Stain Quick kits.

100 mL	TBC-OT-100
250 mL	TBC-OT-250
500 mL	TBC-OT-500
1000 mL	TBC-OT-1L
2500 mL	TBC-OT-2.5L



TB Decolorizer

For use in TB-Stain Hot and TB-Stain Cold kits.

100 mL	TBD-OT-100
100 ML	180-01-100
250 mL	TBD-OT-250
500 mL	TBD-OT-500
1000 mL	TBD-OT-1L
2500 mL	TBD-OT-2.5L



Methylene Blue Loeffler's reagent

Blue counterstain for TB-Stain Hot kit.

100 mL	MBL-OT-100
250 mL	MBL-OT-250
500 mL	MBL-OT-500
1000 mL	MBL-OT-1L
2500 mL	MBL-OT-2.5L





TB-Stain ECO kit

Three-reagent phenol-free kit for staining acid-fast bacteria. Contains TB-Fuchsin reagent, double amount of TB Decolorizer and Methylene Blue Loeffler's reagent as counterstain.

5x100 mL	TBE-K-100
5x250 mL	TBE-K-250
5x500 mL	TBE-K-500



TB-Fuchsin reagent

Primary stain **phenol-free** reagent for acid-fast bacteria used in TB-Stain ECO kit.

100 mL	TBFR-OT-100
250 mL	TBFR-OT-250
500 mL	TBFR-OT-500
1000 mL	TBFR-OT-1L
2500 mL	TBFR-OT-2.5L



Additional TB-Stain Hot kit staining solution

Methylene Blue 10x concentrated

Concentrated counterstain solution of Methylene Blue dye for use in microbiology.

For preparation of 1000 mL of Methylene Blue counterstain solution	100 mL	MB10X-OT-100
For preparation of 5000 mL of Methylene Blue counterstain solution	500 mL	MB10X-OT-500
For preparation of 10000 mL of Methylene Blue counterstain solution	1000 mL	MB10X-OT-1L



TB-Stain Cold and Quick kits

Kits for staining acid-fast bacteria according to Kinyoun and Kinyoun-Gabbett

TB-Stain Quick kit

Three-reagent kit for rapid staining of acid-fast bacteria using Kinyoun-Gabbett method. Contains TB Carbol Fuchsin reagent and TB Armand reagent as counterstain.

3x100 mL	TBQ-K-100
3x250 mL	TBQ-K-250
3x500 mL	TBQ-K-500

TB Armand reagent

Blue counterstain for use in TB Stain Quick kit.

100 mL	TBAR-OT-100
250 mL	TBAR-OT-250
500 mL	TBAR-OT-500
1000 mL	TBAR-OT-1L
2500 mL	TBAR-OT-2.5L

Additional TB-Stain Cold kit staining solution

Gabbett reagent

Blue counterstain for staining acid-fast bacteria according to modified Kinyoun method.

100 mL	GAB-OT-100
250 mL	GAB-OT-250
500 mL	GAB-OT-500
1000 mL	GAB-OT-1L

TB-Stain Cold kit

Three-reagent kit for staining acid-fast bacteria according to Kinyoun. Contains TB Carbol Fuchsin reagent, double amount of TB Decolorizer and TB Malachite Green reagent as counterstain.

for 100 tests	TBC-100T
4x100 mL	TBC-K-100
4x250 mL	TBC-K-250
4x500 mL	TBC-K-500

TB Carbol Fuchsin reagent

Synonym: Ziehl-Neelsen reagent. Primary stain reagent for acid-fast bacteria used in TB-Stain Hot, TB-Stain Cold and TB-Stain Quick kits.

TBC-OT-100
TBC-OT-250
TBC-OT-500
TBC-OT-1L
TBC-OT-2.5L



TB Decolorizer

For use in TB-Stain Hot and TB-Stain Cold kits.

100 mL	TBD-OT-100
250 mL	TBD-OT-250
500 mL	TBD-OT-500
1000 mL	TBD-OT-1L
2500 mL	TBD-OT-2.5L



TB Malachite Green reagent

Green counterstain for TB-Stain Cold kit.

100 mL	TBM-OT-100
250 mL	TBM-OT-250
500 mL	TBM-OT-500
1000 mL	TBM-OT-1L
2500 mL	TBM-OT-2.5L



TB-Stain Fluorescent and TB-Stain Auramine O kits

Kits for fluorescence microscopy detection of acid-fast bacteria according to modified Truant methods

TB-Stain Fluorescent kit

Three-reagent kit for fluorescence microscopy detection of acid-fast bacteria. Contains TB Auramine-Rhodamine reagent, double amount of TB Decolorizer Fluorescent and TB Permanganate reagent as counterstain.

for 100 tests TBF-100T 4x100 mL TBF-K-100 4x250 mL TBF-K-250 4x500 mL TBF-K-500

TB-Stain Auramine O kit

Three-reagent kit for staining acid-fast bacteria using fluorescence method. Contains TB Auramine O reagent, double amount of TB Decolorizer Fluorescent and counterstain of TB Permanganate reagent.

 4x100 mL
 TBAO-K-100

 4x250 mL
 TBAO-K-250

 4x500 mL
 TBAO-K-500

TB Auramine O reagent

Primary stain for use in TB Stain Auramine O kit.

 100 mL
 TBAO-OT-100

 250 mL
 TBAO-OT-250

 500 mL
 TBAO-OT-500

 1000 mL
 TBAO-OT-1L

 2500 mL
 TBAO-OT-2.5L

TB Auramine-Rhodamine reagent

Primary stain reagent for use in TB-Stain Fluorescent kit.

 100 mL
 TBA-OT-100

 250 mL
 TBA-OT-250

 500 mL
 TBA-OT-500

 1000 mL
 TBA-OT-1L

 2500 mL
 TBA-OT-2.5L

TB Decolorizer Fluorescent

For use in TB-Stain Fluorescent and TB-Stain Auramine O kits.

 100 mL
 TBF-OT-100

 250 mL
 TBF-OT-250

 500 mL
 TBF-OT-500

 1000 mL
 TBF-OT-1L

 2500 mL
 TBF-OT-2.5L

TB Permanganate reagent

Counterstain for use in TB-Stain Fluorescent kit and TB-Stain Auramine O kits.

 100 mL
 TBP-OT-100

 250 mL
 TBP-OT-250

 500 mL
 TBP-OT-500

 1000 mL
 TBP-OT-1L

 2500 mL
 TBP-OT-2.5L







Solutions for staining and analysis of fungi

In order to detect and achieve better visualization of fungi and fungal structures that are otherwise difficult to observe, fungi can be stained using Lactophenol solutions. Using Myco-Mount medium they can also be stained and preserved as a permanent preparation.

Lactophenol Blue solution

For use in mycology for staining and analysis of fungi.

100 mL LPB-OT-100 250 mL LPB-OT-250

Lactophenol solution

Colorless solution for use in mycology and analysis of fungi.

100 mL LP-OT-100 250 mL LP-OT-250

For detection and fungal staining of histology sections, please see page 46 for more details on Grocott kit.

Myco-Mount Blue

Medium for simultaneous staining of fungi blue and preparation of permanent slides.

30 mL MYMB-30 100 mL MYMB-100

Myco-Mount Red

Medium for simultaneous staining of fungi red and preparation of permanent slides.

30 mL MYMR-30 100 mL MYMR-100

Potassium hydroxide, solution for fungi

Solution for use in mycology for analysis and visualization of fungi.

30 mL KHOG-OT-30 100 mL KHOG-OT-100



Special staining kits

High quality, efficiency, clearly distinguished tissue components and maximum usability of reagents - these all are characteristics of BioGnost's special staining kits. The kits are easy to use and adapted to fit the user's needs.

Classification of special staining kits according to use

MUSCLE AND/OR CONNECTIVE TISSUE

A.F.O.G kit Masson-Goldner Trichrome kit

Azan Trichrome kit Movat kit
Elastica-Van Gieson kit Orcein kit

Gomori Trichrome kit

H.B.F.P. kit

Paraldehyde Fuchsin kit

P.A.S.M. / Jones kit

Hematoxylin P.T.A. kit Reticulin kit

Hematoxylin W kit Reticulin Contrast kit

Mallory Trichrome kit Van Gieson Trichrome kit

Martius Scarlet Blue (MSB) kit

Verhoeff kit

Masson Trichrome kit Weigert-Van Gieson kit

FUNGI, BACTERIA AND

PARASITES

Alcian Yellow Toluidine Blue kit

BioGram Histo kit

Field kit

Giemsa HP kit

Grocott kit

M.I.F. kit

TB-Stain Histo kit Warthin Starry kit

vvar tilli Starry

LIPIDS

Oil Red O kit

Sudan Black B Lipid kit

NUCLEI AND NUCLEIC ACIDS

Feulgen kit

PIGMENTS AND MINERAL

DEPOSITS

Fouchet-Van Gieson kit

Grimelius kit

HemoGnost Perls kit Masson Fontana kit

Orcein kit Rhodanine kit Von Kossa kit

CYTOLOGY

Bio-Diff RTU kit

Eosin and nigrosin vital kit

Sperm-Diff RTU kit Sudan Black B kit

UriGnost S kit

UriGnost SM kit

CARBOHYDRATES

Alcian Blue - P.A.S. kit Alcian Blue pH 1.0 kit

Alcian Blue pH 2.5 kit

Colloidal Iron kit

Congo Red Highman kit

Congo Red Puchtler kit

Mucicarmine kit

P.A.S. kit

CRYOSTAT

Gomori Trichrome kit


Oil Red O kit

Orcein kit

Sudan Black B Lipid kit

CENTRAL NERVOUS SYSTEM


Luxol Fast Blue kit



A.F.O.G. kit

Six-reagent \underline{A} cid \underline{E} uchsin \underline{O} range \underline{G} kit for selective staining of glomerular protein deposits and collagen in kidney biopsies. Nuclear stain is obtained with Weigert ferric hematoxylin, cytoplasm with Orange \underline{G} and highly selective collagen stain with Aniline blue.


for 100 tests AFOG-100T 6x100 mL AFOG-K-100



Alcian Blue - P.A.S. kit

Seven-reagent Alcian Blue - Periodic Acid-Schiff kit for staining acid mucopolysaccharides according to Mowry. Enables differentiation between acid mucins (stained light blue) and neutral mucins, glycogens and glycoproteins (stained magenta).


for 100 tests AB-100T 7x100 mL AB-K-100 7x500 mL AB-K-500



Alcian Blue pH 1.0 kit

Three reagent kit for specific staining of heavily sulfated muccopolysaccharides blue. Slides are counterstained with Nuclear Fast Red (Kernechtrot) reagent to fully detect the presence of Alcian Blue positive staining.


for 100 tests AB10-100T 3x100 mL AB10-K-100



Alcian Blue pH 2.5 kit

Three reagent kit for staining sulfated and carboxylated acid muccopolysaccharides and sulfated and carboxylated sialomucins blue. Slides are counterstained with Nuclear Fast Red (Kernechtrot) reagent to fully detect the presence of Alcian Blue positive staining.


for 100 tests AB25-100T 3x100 mL AB25-K-100



Alcian Yellow Toluidine Blue kit

Six-reagent kit for staining *Helicobacter pylori* in gastric tissue sections. This method is one of the most popular non-silver method for staining of *H. pylori*, where bacteria are stained blue in contrast to yellow mucins.


for 100 tests AYTB-100T



Azan Trichrome kit

Five-reagent kit for connective tissue staining according to Mallory. Used for visualization of muscle fibers, collagen, glial cells, kidney glomerular stroma, erythrocytes, reticulum and nuclear chromatin in histologic sections.

for 100 tests AZT-100T 5x250 mL AZT-K-250




Bio-Diff RTU kit

Ready-to-Use three-reagent kit with reagents stored in containers that can be used as staining jars. Kit contains fixative reagent, red and blue components for fast and effective staining and buffer tablet for consistent staining results.

3x100 mL

BD-RTU-100






BioGram Histo kit

Five-reagent kit for identification of bacteria according to Gram. For differentiation between Gram-positive and Gram-negative bacteria in histology sections.

for 100 tests


BGRH-100T



Colloidal Iron kit

Six-reagent kit used for visualiziation of carboxylated and sulphated groups of acid mucopolysaccharides and proteogycans. This method can be combined with the P.A.S. method; that way glycogen and neutral mucopolysaccharides would get differentially stained characteristically magenta.

for 100 tests KOL-100T




Congo Red Highman kit

Three-reagent kit for staining amyloids, characteristic for use of alkaline solution as differentiation medium in order to avoid undesirable non-specific coloration of cellular substances. Amyloid deposits display apple green birefringence with polarized light.


for 100 tests 3x100 mL CR-100T CR-K-100





Congo Red Puchtler kit


Three-reagent kit for staining amyloids, characteristic by its high ionic strength and pH enhancing specificity of Congo Red dye binding to amyloid clusters. Method developed by Puchtler remains the gold standard for amyloids in tissue sections. Amyloid clusters have the property of double refraction that enables green coloration under polarized light.



Elastica-Van Gieson kit

Four-reagent kit for staining elastic fibers and differentiation between elastic tissue, collagen and other types of connective tissue. The rapid method enables a satisfactory result with shorter section staining time.


for 100 tests WVGB-100T 4x100 mL WVGB-K-100



Eosin and nigrosine vital kit

Two-reagent kit containing separate dyes for rapid detection of sperm vitality and simple visualization of dead and living sperm cells. The Nigrosin stain provides dark background for easier recognition of both viable and non-viable spermatozoa.


2x30 mL ENVK-K-30 2x100 mL ENVK-K-100



Feulgen kit

Five-reagent DNA staining kit according to Feulgen. For use in semiquantitative DNA determination in histological and cytological samples. Specimen is first treated with hydrochloric acid creating an aldehyde group of DNA that can be visualized by Schiff (BioSchiff) reagent. This reaction is specific for nuclear DNA.


for 100 tests FE-100T 5x100 mL FE-K-100



Field kit

Ready to use two-reagent kit for rapid and efficient staining and detection of parasites in hematology samples. Primarily used for staining thin and thick blood smears (dense drop) for purpose of diagnosing blood parasites. Reagents are stored in containers that can be used as staining jars.


for 100 tests FD-100T



Fouchet-Van Gieson kit

Three-reagent kit for visualization of bilirubin and collagen according to Kutllick. Bilirubin is a yellow-brown pigment, but changes to green due to oxidation induced by Fouchet reaction. Green bilirubin can easily be detected on yellowish and pink colored background.


for 100 tests FVG-100T 3x100 mL FVG-K-100



Giemsa HP kit

Four-reagent kit for staining *Helicobacter pylori* in gastroscopic sections according to Lennart. Advantages of this method for detecting *H. pylori* are sensitive and reproducible results and easy performance.

for 100 tests GMHP-100T




Gomori Trichrome kit

Five-reagent kit for staining muscle, collagen fibre and nuclei, contains blue counterstain. Kit can be used to contrast skeletal, cardiac or smooth muscle.

for 100 tests GT-100T 5x100 mL GT-K-100


 $\ensuremath{^{\star}}$ Gomori Trichrome kit with green counterstain also available



Grimelius kit

Five-reagent kit for staining argyrophilic granules. Grimelius kit can be used for the detection of secretory intracytoplasmatic granules specific for carcinoid tumors and for identification of neuroendocrine cells.


for 100 tests GRI-100T 5x100 mL GRI-K-100



Grocott kit

Six-reagent kit for visualization of fungi and histological argentaffin structures in general (such as basal membranes). Green counterstain provides clear and visually rich contrast to target structures stained black.


for 100 tests GRC-100T 5x100 mL + 3x250 mL GRC-K-100



H.B.F.P. kit

Three-reagent Hematoxylin-Basic Fuchsin-Picric acid staining kit for detection of cardiac muscle changes after ischemia or myocardial infarction. H.B.F.P. kit is a non-enzymatic histochemical technique for detection of early myocardial ischemia with vivid contrast.


for 100 tests HBFP-100T 3x100 mL HBFP-K-100



Hematoxylin P.T.A. kit

Four-reagent Hematoxylin-Phosphotungstic Acid kit for differentiation of smooth and striated muscle tissues as well as for detection of fibrin, collagen and elements of central nervous system according to Mallory.

for 100 tests HPTA-100T 4x100 mL HPTA-K-100



Hematoxylin W kit


Acid-resistant hematoxylin according to Weigert. Two-reagent kit that stains the nuclei blue-black, often a component of special staining kits for connective tissues.

for 100 tests HEMW-100T 2x100 mL HEMW-K-100 2x500 mL HEMW-K-500 2x1000 mL HEMW-K-1L



Three-reagent HemoGnost Perls (Prussian blue / Berlin blue) kit for the detection of reactive ferric (Fe³+) ions in cells. It is often applied on bone marrow and spleen cells.

for 100 tests HP-100T 3x250 mL HP-K-250



Luxol Fast Blue kit

Three-reagent kit for staining myelin and myelinated axons, Nissel bodies and phospholipids according to Kluver-Barrera. Kit is used for identifying the basic neuronal structure in brain or spinal cord sections.


for 100 tests LFB-100T 3x100 mL LFB-K-100



M.I.F. kit

Two-reagent Merthiolate-Iodine-Eormalin kit for fixing and staining fecal parasites (especially protozoae, cysts, helminthic eggs and larvae). Stool sample is fixed with formalin and stained with two coloring agents; iodine and Eosin Y.


for 100 tests MIF-100T



Mallory Trichrome kit

Three-reagent staining kit for connective tissue visualization and detection of collagen, cartilage, muscle, elastic fibers, mucous, pituarity cells, reticulum, bones, amyloid and erythrocytes.

for 100 tests MT-100T 3x100 mL MT-K-100



Martius Scarlet Blue (MSB) kit

Seven-reagent kit used for fibrin visualization, especially of older clusters. This method is a modification of Masson Trichrome and is ideal for studying connective tissue and vascular pathology.


for 100 tests MSB-100T 6x100 mL + 1x250 mL MSB-K-100



Masson Fontana kit

Six-reagent melanine and argentaffin granule staining kit, based on the reduction of silver nitrate to elemental silver. Melanin is a brown-black pigment normally present in the hair, skin, retina, iris and certain parts of CNS. Argentaffin granules are found in carcinoid tumors.


for 100 tests MF-100T 9x100 mL MF-K-100



Masson-Goldner Trichrome kit

Seven-reagent kit for staining muscle and collagen fibers with green counterstain. It is also used for visualizing gametes, nuclei, neurofibrils, glial cells, keratins, intercellular fibrils and for differentiation of smooth muscle fibers and collagens.


for 100 tests MGT-100T 7x100 mL MGT-K-100 7x500 mL MGT-K-500



blue counterstain. It is also used for visualizing gametes, nuclei, neurofibrils, glial cells, keratins and intercellular fibrils. Kit may be useful for the detection of collagen from cancer smooth muscle or diseases like cirrhosis.

for 100 tests MST-100T 7x100 mL MST-K-100 7x500 mL MST-K-500






Movat kit

Ten-reagent kit for staining collagen, elastic and muscle fibers, mucin and fibrin in tissue section. Movat kit is particularly useful when examining heart and vascular diseases.

for 100 tests MOV-100T 10x100 mL MOV-K-100




Mucicarmine kit

Mucicarmine kit is often used to identify primary tumor sites, distinguishing mucin-negative undifferentiated squamous cell lesions from mucin-positive adenocarcinomas. It can also be used as indicative of diseases such as asthma, bronchitis, chronic obstructive pulmonary disease and cystic fibrosis.

MUC-100T for 100 tests 4x100 mL MUC-K-100






Oil Red O kit

Four-reagent kit for selective staining and detection of fat cells and neutral fats according to Johnson. It can be used with frozen sections and fresh smears to detect obesity-linked pathologies such as dyslipidemia and diabetes type.

ORO-100T for 100 tests 4x250 mL ORO-K-250




Orcein kit

Five-reagent kit for visualization of hepatitis B surface antigen (HBsAg) seen as viral inclusion bodies in hepatocytes, for elastic fibers and copper associated protein in tissue sections. It can be used with frozen sections.

for 100 tests ORC-100T 5x250 mL ORC-K-250






Paraldehyde Fuchsin kit

Seven-reagent kit according to Gomori for detecting pathological changes in elastic fibers. It also stains mast cell granules, beta granules in pancreatic islets, neurosecretory material, mast cell granules and beta cells in the pituitary gland.


for 100 tests PAF-100T 9x100 mL PAF-K-100



P.A.S. kit

Five-reagent Periodic Acid-Schiff kit for staining aldehydes, muccopolysaccharides, mucoproteins and lymphocytes according to Hotchkiss-McManus. P.A.S. staining may also be used for the demonstration of fungal organisms in tissue sections.


for 100 tests PAS5-100T 5x100 mL PAS5-K-100 5x500 mL PAS5-K-500



P.A.S.M. / Jones kit

Six-reagent <u>Periodic Acid Silver Methenamine kit for staining</u> kidney glomerular basement membranes. Kit includes red counterstain which provides visually rich contrast to target structures stained black.


for 100 tests PASM-100T 5x100 mL + 3x250 mL PASM-K-100



Reticulin kit

Seven-reagent kit for detecting argyrophilic reticulin fibers. It clearly differentiates between collagen and reticulin and nerve fibers and connective tissue. The main function of reticular fibers is to provide support and are normally found in liver, lymph nodes, spleen and kidneys.

for 100 tests RE-100T 7x50 mL RE-K-50 7x100 mL RE-K-100



Reticulin Contrast kit

Nine-reagent kit for detecting argyrophilic reticulin fibers according to Gordon and Sweets. The kit contains gold chloride solution that enhances visualization of reticulin fibers and it also contains Nuclear Fast Red (Kernechtrot) reagent that enables fine contrasting background.

for 100 tests RET-100T 9x50 mL RET-K-50 9x100 mL RET-K-100






Rhodanine kit

Four-reagent kit for detecting copper and copper-associated protein (CAP) in patients suffering from Wilson's disease. Abnormal copper accumulations are predominately found in liver tissue, but can also be found in the brain and cornea of the eyes.

for 100 tests ROD-100T 5x100 mL ROD-K-100




Sperm-Diff RTU kit

Ready-to-Use three-reagent kit with reagents stored in containers that can be used as staining jars. Kit contains fixative reagent, red and blue components for fast and effective staining.

3x100 mL SP-RTU-100

49



Sudan Black B kit

Four-reagent kit for staining neutrophil granules in hematological smears (blood or bone marrow films). Used as one of the methods for detecting myelocytic and myelomonocytic leukemia.


for 350-400 tests SBBH-K-100



Sudan Black B Lipid kit

Four-reagent kit for specific lipid staining in cytochemistry. Sudan Black dye stains a few types of lipids, including neutral fats, phospholipids and sterols. Contains double amount of Sudan Black B, solution.


for 100 tests SBB-100T 5x100 mL SBB-K-100



TB-Stain Histo kit

Three-reagent kit for staining acid-fast bacteria (pathogenic mycobacteria) in histology sections, sputum, smears and culture smears according to Ziehl-Neelsen. Heating of the carbol-fuchsin solution is avoided in this protocol hence the release of hazardous phenolic vapors is omitted.

for 100 tests TBHK-100T



UriGnost S kit

Kit for sampling, staining, and microscopic analysis of urine sediment. Contains modified reagent according to Sternheimer (reagent manufactured according to the European Confederation of Laboratory Medicine (ECLM) and all the necessary equipment for sampling, concentrating, staining, and counting cells and kidney cylinders, as well as urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 μL pipette tips and urine plates).


for 500 tests USK-500 * for more information, see pages 57 and 75



UriGnost SM kit

Kit for sampling, staining and microscopic analysis of urine sediments. Contains modified reagent according to Sternheimer-Malbin and equipment required for sampling, concentration, staining, counting of kidney cells and kidney cylinders and urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 μ L pipette tips and urine plates).


for 500 tests USMK-500 * for more information, see pages 57 and 76



Van Gieson Trichrome kit

Three-reagent kit for staining collagen fibers, muscle tissue, keratinized epithelium, cytoplasm, glial fibers and erythrocytes. Used for differentiation between collagen and smooth fibers in tumors and various other diseases.


for 100 tests VG-100T 3x100 mL VG-K-100



Verhoeff kit

Six-reagent kit for detecting atrophy of elastic tissue in cases of emphysema, thinning and loss of elastic fibers in arteriosclerosis and other vascular diseases, or whether blood vessels have been invaded by a tumor.

for 100 tests VER-100T 6x100 mL VER-K-100




Von Kossa kit

Five-reagent kit for simple and reproducible detection of calcium deposits and calcium salt in tissue samples according to Von Kossa. Tissue calcification is associated with metabolic problems in various tissues (like bone marrow and mamma) and in tumors.

for 100 tests VK-100T 5x100 mL VK-K-100






Warthin Starry kit

Five-reagent kit for staining *Spirochaeta*, *Helicobacter pylori*, *Microsporadia* and *Legionella pneumophila*. The kit contains 12 jars with gelatin that enables both incubation and staining of sections, as well as other reagents that enable precipitation of silver on the bacterial surface. The bacteria are found in the mucus of the surface epithelium, in the apical gastric glands and in the gastric mucosa.

up to 48 tests WS-12T





Weigert-Van Gieson kit

Six-reagent kit for staining elastic fibers with prolonged incubation period. Used for differentiation between elastic tissue, collagen and other types of connective tissue.

for 100 tests WVG-100T 6x100 mL WVG-K-100











Staining reagents and solutions

10% Formalin, alcoholic solution

For use with Reticulin and Reticulin Contrast staining kits.

100 mL F4A-OT-100

Acetic acid, 0.5% solution

For use with special kits.

100 mL OKO5-OT-100

Acid alcohols

For use with Azan Trichrome and Orcein staining kits.

 Acid alcohol, Azan
 100 mL
 KAA-OT-100

 250 mL
 KAA-OT-250

 Acid alcohol, Orcein
 250 mL
 KAO-OT-250

Acid solution

For preparation of impregnation solution with Warthin Starry kit.

500 mL KOT-OT-500

Acridine Orange solution

Acridine Orange aqueous solution is used as fluorescent dye for differential staining of DNA and RNA, acid mucins and detecting apoptosis; in cytogenetics for displaying structures containing abundant amounts of DNA. Also enables detection of microorganisms in blood smears.

 100 mL
 AOA-OT-100

 500 mL
 AOA-OT-500

 1000 mL
 AOA-OT-1L

A.F.O.G. reagent

For use with A.F.O.G. (Acid Fuchsin Orange G) kit.

100 mL AFOG-OT-100

Alcian Blue solution 1%

Alcian Blue 8GX dye 1% aqueous solution.

100 mL AB1-OT-100 500 mL AB1-OT-500

Alcian Blue solution pH 1.0

Reagent for staining heavily sulfated mucopolysaccharides. A component of Alcian Blue pH 1.0 kit.

100 mL AB10-OT-100 500 mL AB10-OT-500 1000 mL AB10-OT-1L

Alcian Blue solution pH 2.5

Reagent for staining acid mucopolysaccharides of intestinal goblet cells. A component of Alcian Blue-P.A.S. and Alcian Blue pH 2.5 staining kits.

100 mL AB2-OT-100 500 mL AB2-OT-500 1000 mL AB2-OT-1L

Alkaline alcohol, solution

For use with MSB kit.

100 mL ALA-OT-100

Ammonium iron sulfate, solution

For use with Reticulin and Reticulin Contrast staining kits.

100 mL ASF-OT-100

Aniline, alcoholic solution

For use with Azan Trichrome kit.

100 mL ANA-OT-100 250 mL ANA-OT-250

Aniline Blue reagent

Counterstain used with Masson Trichrome kit.

100 mL ABR-OT-100 500 mL ABR-OT-500

Aniline Blue, solution

For use with MSB kit.

100 mL ABO-OT-100

Azan reagent

For use with Azan Trichrome kit.

100 mL AZR-OT-100 250 mL AZR-OT-250

Azocarmine, solution

For use with Azan Trichrome kit.

100 mL AZ-OT-100 250 mL AZ-OT-250

Basic Activation buffer

For use with Oil Red O kit.

250 mL BAP-OT-250

Benedict's reagent

For detecting glucose in urine.

 100 mL
 BNR-OT-100

 250 mL
 BNR-OT-250

 500 mL
 BNR-OT-500

 1000 mL
 BNR-OT-1L

Biebrich Scarlet-Acid Fuchsin reagent

For use with special kits.

100 mL BSAF-OT-100 500 mL BSAF-OT-500

BioSchiff reagent

Schiff's reagent for detecting aldehydes and mucous substances. A component of P.A.S. and Alcian Blue-P.A.S. staining kits.

100 mL BS-OT-100 500 mL BS-OT-500 1000 mL BS-OT-1L

Borax, solution

For use with Grocott and P.A.S.M./Jones kits.

100 mL BO-OT-100 250 mL BO-OT-250

Chromotrope 2R / Aniline Blue reagent

Blue counterstain used with Gomori Trichrome kit.

100 mL CABR-OT-100

Chromotrope 2R / Fast Green F.C.F. reagent

Green counterstain used with Gomori Trichrome kit.

100 mL CFG-OT-100

Colloidal Iron, stock solution

Reagent for binding ferric ions to acid mucopolysaccharides and proteoglycans.

100 mL KOL-OT-100

Congo Red Puchtler reagent

For use with Congo Red Puchtler kit.

100 mL CRPR-OT-100

Congo Red reagent

For use with Congo Red Highman kit.

100 mL CR-OT-100

Cresyl Violet, solution

For use with Luxol Fast Blue kit.

100 mL CV-OT-100

Differentiation reagents based on acetic acid

For use with special staining kits in histopathology.

Reagent for differentiation in Gomori Trichrome kit	100 mL	RDG-OT-100
Acetic acid, 1% solution (used with Masson Trichrome and Masson-Goldner staining kits)	100 mL 500 mL	OK1-OT-100 OK1-OT-500
Acetic acid, Colloidal Iron (A) solution	250 mL	OKA-OT-250
Acetic acid, Colloidal Iron (B) solution	100 mL	OKB-OT-100

Differentiation reagents based on hydrochloric acid

For use with special staining kits in histopathology.

HCL reagent, A.F.O.G. • HCL reagent, Colloidal Iron • HCL reagent, Feulgen • HCL reagent, HemoGnost Perls HCL reagent, HemoGnost Perls HCL reagent, P.A.S.	100 mL 100 mL 100 mL 100 mL 250 mL 100 mL	HCLAF-OT-100 HCLK-OT-100 HCLF-OT-100 HCLH-OT-100 HCLH-OT-250 HCLP-OT-100
HCL reagent, P.A.S. HCL reagent, P.A.S.	100 mL 500 mL	HCLP-OT-100 HCLP-OT-500
HCL reagent, Weigert-Van	100 mL	HCLW-OT-100
Gieson •		

Ehrlich aldehyde reagent

Reagent for detecting urobilinogen in urine.

100 mL EAR-OT-100

Eosin-nigrosin vital

Fast detection (one-step detection) of sperm vitality and visualization of dead and living sperm cells with one reagent. Simple, easy and fast method for semen analysis.

30 mL ENV-30 100 mL ENV-100



Staining reagents and solutions

Eosin Y, isotonic solution

Aqueous solution of Eosin Y used with Eosin and nigrosin vital kit.

100 mL EIO-OT-100

Fast Green F.C.F. reagent

Counterstain used with Masson-Goldner Trichrome and Paraldehyde Fuchsin staining kits.

100 mL FGR-OT-100 500 mL FGR-OT-500

Fast Green F.C.F. contrast reagent

Counterstain used with Grocott staining kit.

100 mL FGKR-OT-100

Field, solution A

Blue component of Field kit.

100 mL FDA-RTU

Field, solution B

Red component of Field kit.

100 mL FDB-RTU

Ferri reagent, Verhoeff B

For use with Verhoeff kit.

100 mL FRV-OT-100

Ferri reagent, Weigert B

Iron(III) nitrate solution in diluted hydrochloric acid, a component of Hematoxylin W kit.

 100 mL
 FR-OT-100

 500 mL
 FR-OT-500

 1000 mL
 FR-OT-1L

Fuchsin Acid reagent

For use with Mallory Trichrome kit.

100 mL FA-OT-100

Fuchsin Acid Van Gieson reagent

For staining with Van Gieson Trichrome, Weigert-Van Gieson and Colloidal Iron kits. Synonym: Picrofuchsin reagent.

100 mL FAG-OT-100

Fuchsin Basic reagent

For use in bacteriology.

 100 mL
 FBR-OT-100

 500 mL
 FBR-OT-500

 1000 mL
 FBR-OT-1L

Fuchsin Basic, solution

For use with H.B.F.P. kit.

100 mL FBO-OT-100

Gelatin, gel in transport jar

For preparing developing solution with Warthin Starry staining kit.

12x8 mL GEL-128

Glutaraldehyde, 4% buffered solution

Solution used with Sudan Black B kit.

100 mL G4-OT-100

0

Gold chloride solutions

Aqueous solutions of gold chloride (aurichloric acid, HAuCl₄), component od many special staining kits.

Gold chloride, 0.2% solution 100 mL ZK02-OT-100 Gold chloride, 1% solution 50 mL ZK1-OT-50 Gold chloride, 2% solution 50 mL ZK2-OT-50

Hematoxylin, Verhoeff A

For use with Verhoeff kit.

100 mL HEMV-OT-100

Hematoxylin, Weigert A

Hematoxylin solution in ethanol, component of Hematoxylin W kit.

 100 mL
 HEMA-OT-100

 500 mL
 HEMA-OT-500

 1000 mL
 HEMA-OT-1L

Hydroquinone, solution

For preparation of developing solution in Warthin Starry kit.

50 mL HQ-OT-50

Iodine solution, Verhoeff C

For use with Verhoeff kit.

100 mL JODV-OT-100

Iron chloride, solution

For use with Fouchet-Van Gieson kit.

100 mL FK-OT-100

Iron nitrate, solution

Iron nitrate solution for use with Movat kit.

100 mL FN-OT-100

Kovac indole reagent

Reagent for identification of indole-positive and indole-

negative microorganisms.

100 mL KOV-OT-100

Light Green S.F. reagent

Counterstain used with trichrome special staining kits.

100 mL LGS-OT-100

Lithium carbonate, Luxol

Lithium carbonate solution used with Luxol Fast Blue kit.

100 mL LKL-OT-100

Lithium carbonate, solution

Lithium carbonate solution used with Von-Kossa staining kit.

100 mL LK-OT-100

Lugol's solution, M.I.F.

For use with M.I.F. kit.

100 mL LUGM-OT-100

Luxol Fast Blue, solution

For use with Luxol Fast Blue kit.

100 mL LFB-OT-100

M.F. reagent

For use with M.I.F. kit.

100 mL MF-OT-100

Martius Yellow, solution

For use with MSB kit.

100 mL MAY-OT-100

Mayer's mucicarmine

For use with Mucicarmine kit.

100 mL MMU-OT-100

Metanil Yellow, solution

For use with Mucicarmine kit.

100 mL MY-OT-100

Methenamine, solution

For use with Grocott and P.A.S.M./Jones kits.

100 mL MET-OT-100 250 mL MET-OT-250

Methyl Blue Eosin, solution

Modified Mann's solution for use in histology.

 100 mL
 MBE-OT-100

 500 mL
 MBE-OT-500

 1000 mL
 MBE-OT-1L

Neutral Red reagent

Neutral Red aqueous counterstain solution for use with special staining kits.

100 mL NR-OT-100

Nigrosine, isotonic solution

Aqueous solution of nigrosine used with Eosin and nigrosine vital kit.

100 mL NIO-OT-100

Nuclear Fast Red (Kernechtrot) reagent

Nuclear Fast Red dye solution for staining nuclei. Counterstain in many special staining kits.

 100 mL
 KR-OT-100

 250 mL
 KR-OT-250

 500 mL
 KR-OT-500

Oil Red O reagent

Lipid substances staining solution.

250 mL ORO-OT-250

Orange G, 1% solution

For use with Movat kit.

100 mL ORG1-OT-100

Orange G / Aniline Blue reagent

For use with Mallory Trichrome kit.

100 mL OGA-OT-100

Orcein reagent

For use with Orcein kit.

250 mL ORCR-OT-250

•

Oxalic acid solutions

For use with Hematoxylin P.T.A., Masson Fontana, Orcein, Paraldehyde Fuchsin and Reticulin staining kits.

Oxalic acid, 1% solution	100 mL	OKS1-OT-100
Oxalic acid, 2% solution	250 mL	OKS2-OT-250

Paraldehyde Fuchsin reagent

Reagent for staining elastins and mucins.

100 mL PAF-OT-100



Periodic acid, solution

Synonym: Orthoperiodic acid. Oxidizing medium widely used in histology. Key component of many special staining kits, such as P.A.S. and Alcian Blue-P.A.S., P.A.S.M. / Jones, Grocott and Weigert-Van Gieson kit.

Periodic acid, 0.5% solution	100 mL	PK05-OT-100
Periodic acid, 0.5% solution	250 mL	PK05-OT-250
Periodic acid, 0.5% solution	500 mL	PK05-OT-500
Periodic acid, 0.8% solution	100 mL	PK08-OT-100
Periodic acid, 0.8% solution	250 mL	PK08-OT-250
Periodic acid, 0.8% solution	500 mL	PK08-OT-500
Periodic acid, 1% solution Periodic acid, 1% solution Periodic acid, 1% solution	100 mL 250 mL 500 ml	PK1-OT-100 PK1-OT-250 PK1-OT-500

Phosphate buffered saline (PBS)

Sterilized isotonic phosphate buffer solutions.

Buffered saline (Dulbecco), without added calcium and magnesium, sterile filtered	500 mL	D-OT-500
Buffered saline (Dulbecco), without added calcium and magnesium, concentrated 10x, sterile filtered	500 mL	DX-OT-500
Buffered saline (Dulbecco), with added calcium and magnesium, sterile filtered	500 mL	DCM-OT-500
Buffered saline (Dulbecco), with added calcium and magnesium, concentrated 10x, sterile filtered	500 mL	DCMX-OT-500
Buffered saline (Dulbecco) (0.01 mol/L, Tween 20), sterile filtered	500 mL	PBS-OT-500

Staining reagents and solutions

Phosphomolybdic acid, 1% solution

For use with A.F.O.G. (Acid Fuchsin Orange G) and Mallory Trichrome staining kits.

FMK1-OT-100 100 ml

Phosphomolybdic acid, 5% solution

For use with Azan Trichrome kit.

100 mL FMK5-OT-100 250 mL FMK5-OT-250

Phosphotungstic acid, 1% solution

For use with Mallory Trichrome kit.

100 ml FVK1-OT-100 500 mL FVK1-OT-500

Phosphotungstic acid, 5% solution

For use with Movat kit.

100 mL FVK5-OT-100

Picric acid in acetone, solution

For use with H.B.F.P. kit.

100 mL PKA-OT-100

Ponceau S, solution

For use with MSB kit.

100 mL PONS-OT-100

Potassium hexacyanoferrate, Colloidal Iron solution

Potassium hexacyanoferrate (II) solution for use with Colloidal

Iron kit.

100 mL KHCK-OT-100

Potassium hexacyanoferrate, solution

Potassium hexacyanoferrate (II) solution, optimally concentrated for use with HemoGnost Perls kit.

100 mL KHC-OT-100 250 mL KHC-OT-250

Potassium hydroxide, solution

Optimally concentrated aqueous solution of potassium

hydroxide used with Congo Red Highman kit.

100 mL KHO-OT-100

Potassium permanganate, solutions

For use with Hematoxylin P.T.A., Masson Fontana, Orcein, Paraldehyde Fuchsin and Reticulin staining kits.

Potassium permanganate, 100 mL KP05-OT-100

0.5% solution •

Potassium permanganate, 250 mL KP1-OT-250

1% solution •

Propylene glycol

For use with Sudan Black B Lipid kit.

PG-100

Propylene glycol, 85%

For use with Sudan Black B Lipid kit.

PG85-100 100 ml

P.T.A. - P.M.A. reagent

Reagent containing optimal concentration of phosphotungstic and phosphomolybdic acid. Used with Masson Trichrome and Masson-Goldner Trichrome staining kits.

PPR-OT-100 500 mL PPR-OT-500

Reagent for differentiation in Verhoeff kit

For use with Verhoeff kit.

100 mL RDV-OT-100

Reducing buffer, solution

For use with Von Kossa kit.

100 mL RP-OT-100

Resorcine Fuchsin reagent

For use with Weigert-Van Gieson kit.

100 mL RFR-OT-100

Rhodanine reagent

For use with Rhodanine kit.

100 ml RR-OT-100

Silver nitrate solutions

For use in special staining kits.

Silver ammonia solution 100 mL (component of Reticulin and

Reticulin Contrast staining

kits) • •

Silver ammonia reagent (component of Von Kossa and

Masson Fontana staining

kits) • •

Silver nitrate, solution (component of Grocott and

P.A.S.M. / Jones staining

kits) • •

Silver nitrate, WS solution (component of Warthin Starry 100 mL

100 mL

100 mL

SNWS-OT-100

SA-OT-100

SAR-OT-100

SN-OT-100

staining kit)

Sodium acetate, solution

For use with Rhodanine kit.

100 mL NA-OT-100

Sodium hydroxide, solution

For use with Congo Red Puchtler kit.

30 mL NH-OT-30

Sodium metabisulfite solutions

Component of Alcian Blue-P.A.S., P.A.S. and Feulgen staining kits.

Sodium metabisulfite 100 ml NM-OT-100 500 mL NM-OT-500 100 mL NMF-OT-100 Sodium metabisulfite, solution for Feulgen

Sodium tetraborate, solution

A component of Alcian Blue pH 2.5 and Alcian Blue-P.A.S. staining kits.

100 mL NTB-OT-100 500 mL NTB-OT-500

Sodium thiosulfate solution

Solutions are component of Grocott, Masson Fontana, P.A.S.M. / Jones, Reticulin, Reticulin Contrast and Von Kossa staining kits.

Sodium thiosulfate, 2% solution	100 mL	NT2-OT-100
Sodium thiosulfate, 5% solution	100 mL	NT5-OT-100

Sudan Black B, solution

Solution for staining lipids, used with Sudan Black B lipid kit.

100 mL SBO-OT-100

Sudan Black B reagent

Staining reagent for use in hematology, used with Sudan Black B kit.

100 mL SBB-OT-100

Sulphosalicylic acid, 20% solution

Reagent for detecting proteins in urine.

100 mL SSK-OT-100

Sulfuric acid solutions

Used with Orcein, Paraldehyde Fuchsin and Reticulin staining kits.

Sulfuric acid, 0.3% solution	250 mL	SK03-OT-250
Sulfuric acid, 0.5% solution	100 mL	SK05-OT-100
Sulfuric acid, 3% solution	100 mL	SK3-OT-100

Tincture of iodine, 1%

Alcoholic iodine solution.

100 mL TJ-OT-100 1000 mL TJ-OT-1L

Trichloroacetic acid, solution

For use with Fouchet-Van Gieson kit.

100 mL TKO-OT-100

Toluidine Blue, solution

Solution for nuclear staining and metachromatic staining of mastocytes in connective tissue.

100 mL	TB-OT-100
250 mL	TB-OT-250

UriGnost S reagent

Modification according to Sternheimer for staining and microscopical analysis of urine sediment. A component of UriGnost S kit.

UriGnost S reagent for 500 tests	50 mL	UGS-500
UriGnost S reagent for 1000 tests * for more information, see pages	100 mL 50 and 75	UGS-1000

UriGnost SM reagent

Modification according to Sternheimer-Malbin for staining and microscopical analysis of urine sediment. A component of UriGnost SM kit.

UriGnost SM reagent for 500 tests	50 mL	UGSM-500
UriGnost SM reagent for 1000 tests * for more information, see page	100 mL s 50 and 75	UGSM-1000



Note: contact us for information about larger and more affordable packagings (500 g and larger)

Acridine Orange, C.I. 46005

Basic Orange 14. For staining acid mucopolysaccharides, apoptosis detection, differential staining of RNA and DNA.

25 g AO-P-25

Alcian Blue 8 GX, C.I. 74240

Ingrain Blue 1, BSC certified. For staining acid mucopolysaccharides according to Scott, Mowry and Dorling.

5 g	CA8GX-P-5
10 g	CA8GX-P-10
25 a	CA8GX-P-25

Alcian Green

Ingrain Green 2. For staining cartilage, sulfated and carboxylated mucopolysaccharides.

10 q ALG-P-10

Alcian Yellow, C.I. 12840

Ingrain Yellow 1. Used for distinguishing types of glycosaminoglycans and for detection of neurosecretory material.

10 g AY-P-10

Amido Black 10B, C.I. 20470

Naphthol blue black, Acid Black 1. For collagen staining according to Van Gieson trichrome and for hemoglobin and hemosiderin staining according to Puchtler-Swear.

10 g	AMB-P-10
25 g	AMB-P-25

Aniline Blue, C.I. 42780

Methyl Blue, Cotton Blue, Ink Blue, China blue, Acid Blue 93, BSC certified. For staining connective tissue using Mallory and Gomori trichrome methods.

25 g CAB-P-25

Auramine O, C.I. 41000

Basic Yellow 2, BSC certified stain. For staining acid-fast bacteria.

25 g AU-P-25

Azocarmine G, C.I. 50085

Rosinduline. For staining according to AZAN trichrome method according to Heidenhain.

5 g AZC-P-5

Azophloxine, C.I. 18050

Red 2G, Acid red 1. For staining according to Masson trichrome methods (used instead of red Biebrich Scarlet and Fuchsin Acid stains).

5 g	AZO-P-5
25 g	AZO-P-25

Azure A, C.I. 52005

Methylene Azure. For metachromatic staining of mast cells.

Azure B, C.I. 52010

Azure I. For preparation of Romanowsky stains, for staining nuclei or nuclear components.

25 g AZB-P-25

Azure C, C.I. 52002

Azure C blue. For staining cell nuclei and nuclear components, such as DNA.

25 g AC-P-25

Azure II

Azure II. For preparation of Romanowsky stains, made by mixing Azure B and Methylene Blue stains.

5 g A2-P-25

Azure II Eosinate

Azure II eosinate. For preparation of Romanowsky stains, made by mixing equal ratios of Azure B, Methylene Blue and Eosin Y stains.

25 g A2E-P-25

Biebrich Scarlet, C.I. 26905

Croceine Scarlet, Ponceau B, Ponceau BS, BSC certified stain. For Masson trichrome staining and a component of Shorr's solution.

25 g BS-P-25

Bismarck Brown Y, C.I. 21000

Bismark Brown Y, Vesuvine BA, BSC certified. For use in histopathology and cytology.

25 g BB-P-25

Brilliant Cresyl Blue, C.I. 51010

For use in hematology (for reticulocyte staining) and histology (for acid mucopolysaccharides).

10 g BCB-P-10 25 g BCB-P-25

Brilliant Green Certified, C.I. 42040

Malachite Green G, Basic Green 1, Emerald Green. For bacteriostatic use.

25 g BG-P-25

Carmine, C.I. 75470

Natural Red 4, Cochineal, Nacarat, carminic acid and aluminum compound. For staining mucins and glycogens.

5 g CAR-P-5 10 g CAR-P-10 25 g CAR-P-25

Carminic acid, C.I. 75470

Natural Red 4. For staining mucines and glycogen.

5 g CARA-P-5 10 g CARA-P-10 25 g CARA-P-25

Celestine Blue, C.I. 51050

Celestine blue B, Mordant blue 14. For staining nuclei dark blue, used as hematoxylin substitute.

10 g	CB-P-10
25 g	CB-P-25

Chromotrope 2R, C.I. 16570

Acid Red 29, Mordant Blue 80. For counterstaining cytoplasm in histopathology used as eosin substitute in HE staining and in trichrome stains

25 g C2R-P-25

Congo Red, C.I. 22120

Direct Red 28, Kongorot, BSC certified stain. For staining of amyloids according to Highman and Puchtler and for elastic fiber staining.

25 a	CR-P-25

Cresyl Violet Acetate Certified

Cresyl Violet, Oxazine 9, Cresyl Echt Violet. For use in microbiology and histology.

25	g	CVA-P-25

Crystal Violet, C.I. 42555

Gentian Violet, Methyl Violet 10B, BSC certified stain. For bacterial staining according to Gram.

25 g	CRV-P-25
50 g	CRV-P-50

Eosin B, C.I. 45400

Eosin B, Eosin Bluish, Imperial Red, Acid Red 91, BSC certified. For staining according to Romanowsky. Used for staining red blood cells, connective tissues and as counterstain for the Gram stain.

25 g	EB-P-25

Eosin Y, C.I. 45380

Eosin Yellowish, Acid Red 87, Eosin WS, BSC certified stain. Counterstain for hematoxylin and for preparation of EA Pap stains and polychromatic hematological stains.

25 g	EOY-P-25
100 g	EOY-P-100
500 g	EOY-P-500

Eriochrome Cyanine R

Mordant Blue 3, Chromoxane Cyanine R. For use in cytology and histology.

25 g		ECR-P-25
23 y		ECK-F-2.

Fast Green F.C.F., C.I. 42053

Food Green 3, Solid Green F.C.F, 1724 Green, BSC certified stain. For trichrome staining according to Gomori and Masson.

25 g FG-	P-25
----------	------

Fuchsin Acid, C.I. 42685

Acid fuchsine, Acid Violet 19, BSC certified stain. For staining connective tissue according to Van Gieson and Mallory.

25 a	FA-P-25

Fuchsin Basic, C.I. 42510

Basic fuchsine, Basic Violet 14, BSC certified stain. For staining mucins, Negri bodies and acid-fast bacteria.

25 g	FB-P-25
100 g	FB-P-100

Fuchsin Basic - Schiff, C.I. 42500

Pararosaniline HCl, Basic Red 9, BSC certified stain. Used for P.A.S. (Periodic Acid Schiff) staining, DNA staining according to Feulgen and acid-fast bacteria staining.

25 g	FBS-P-25
50 g	FBS-P-50

Gentian Violet, C.I. 42555

Crystal Violet, Methyl Violet 10B, BSC certified stain. For bacterial staining according to Gram.

25 g	GV-P-25
100 a	GV-P-100

Giemsa

BSC certified stain. Romanowsky dye for staining according to Pappenheim. Polychromatic dye used for identification of chromosomal damage and as a differentiating stain.

25 g	G-P-25
50 g	G-P-50

Hematoxylin, C.I. 75290

Natural Black 1, BSC certified stain. For preparation of nuclear staining reagents.

25 g	H-P-25
100 g	H-P-100

Janus Green B, C.I. 11050

Diazine Green, BSC certified stain. For displaying mitochondria in living cells.

25 ~	JG-P-25
25 g	JG-P-25

Jenner Stain

Jenner, BSC certified stain. Romanowsky polychromatic stain routinely used in hematology for staining blood smears.

25 g JS-P-25

Leishman Stain

Eosin-polychrome Methylene Blue. Romanowsky polychromatic dye. Used to distinguish and identify leukocytes and malaria parasites.

25 g LS-P-25	
--------------	--

Light Green S.F., C.I. 42095

Acid Green 5, Pencil Green SF, BSC certified stain. For counterstaining according to Papanicolaou and Masson trichrome methods.

25 g	LG-P-25
------	---------

Luxol Fast Blue MBSN, C.I. 74180

Luxol Fast Blue, Solvent Blue 38. For use in histology.

25 g	LFB-P-25	

Malachite Green, C.I. 42000

Malachite Green Oxalate, Basic Green 4, Victoria Green, BSC certified stain. Used as counterstain in TB-Stain Cold kit.

10 g MAG-P-10 25 g MAG-P-25

Martius Yellow C.I. 10315

Golden Yellow, Martinsgelb, Acid yellow 24. For staining erythrocytes yellow in trichrome staining methods.

25 g MAY-P-25

May-Gruenwald

Polychromatic Romanowsky dye. For blood smears and bone marrow staining.

25 g MG-P-25 50 g MG-P-50

Metanil Yellow, C.I. 13065

Acid yellow 36. For staining cytoplasm, muscle connectors and identifying parietal cells of gastric glands.

25 g MEY-P-25 50 g MEY-P-50 100 g MEY-P-100

Methyl Green, C.I. 42590

For staining nuclei green.

25 g MGR-P-25

Methyl Violet 10B, C.I. 42555

Gentian Violet, Crystal Violet, BSC certified stain. For bacterial staining according to Gram.

25 g MV10B-P-25 50 g MV10B-P-50

Methyl Violet 2B, C.I. 42535

Basic Violet 1, BSC certified stain. For bacterial staining according to Gram.

25 g MV2B-P-25

Methylene Blue, C.I. 52015

Basic Blue 9, BSC certified stain. Used for the preparation of Romanowsky polychromatic stain for staining of bone marrow and as a counterstain in staining of acid-fast bacteria.

25 g MB-P-25 100 g MB-P-100

Methylene Green, C.I. 52020

Methylene Green, BSC certified stain. For staining nuclei green.

25 g MEG-P-25

Neutral Red, C.I. 50040

Basic Red 5, BSC certified stain. For supravital staining of blood cells using Neutral Red-Janus Green B method.

25 g NR-P-25

Nigrosine WS, C.I. 50420

Nigrosine water soluble, Nigrosine B. For use in microbiology and cytology, as negative staining.

25 g NGR-P-25

Nuclear Fast Red, C.I. 60760

Kernechtrot. Used for staining nuclei red by creating aluminum complex.

5 g NF-P-5

Oil Red O, C.I. 26125

Sudan Red 5B, BSC certified stain. For staining lipids.

25 g ORO-P-25

Orange II, C.I. 15510

Acid Orange 7, BSC certified stain. Used as a counterstain, as a substitute for Orange G.

25 q O2-P-25

Orange G, C.I. 16230

Acid Orange 10, Acid Fast Orange G, BSC certified stain. For staining keratin according to Papanicolaou, counterstain for hematoxylin.

25 g OG-P-25 100 g OG-P-100

Orcein, C.I. (ed.1) 1242

Natural Red 28, Orcein Synthetic. For staining elastic tissues and viral inclusion bodies of the hepatitis B surface antigen.

5 g ORC-P-5 10 g ORC-P-10

Phloxine B, C.I. 45410

Acid Red 92, Eosin 10B, Eosin Blue, BSC certified stain. For Methylene Blue-Phloxine staining and staining keratin and mucus according to Kreyberg.

25 g PB-P-25

Ponceau S, C.I. 27195

Acid Red 112, Fast Ponceau 2B. For use in electrophoresis.

25 q PS-P-25

Potassium permanganate

Synonym: Potassium (VII) manganate, Condy's crystals. Potassium permanganate for use in microscopy.

50 g KPH-P-50 100 g KPH-P-100 500 g KPH-P-500

Pyronine B, C.I. 45010

Pyronine B, BSC certified stain. For bacterial, mold and RNA staining.

25 g PYB-P-25

Resorcinol, C.I. 76505

Resorcin, 1.3-benzendiol. Used for differentiation between aldoses and ketoses in Seliwanoff's test by staining ketose sugars dark red.

25 g RES-P-25

Rhodamine B, C.I. 45170

Basic Violet, Acid Red 52. For staining acid-fast bacteria.

25 g RHB-P-25

Rhodanine

DMAB-Rhodanine. For staining deposits of copper bound to proteins.

5 g RHO-P-5

Rose Bengal, C.I. 45440

Acid Red 94, BSC certified stain. For staining in microscopy and histology.

25 g RB-P-25

Safranin O, C.I. 50240

Basic Red 2, Safranine T, Phenazinium, BSC certified stain. Counterstain in the Gram method for detecting Gram-negative bacteria.

25 g SO-P-25

Sirius Red, C.I. 35780

Direct Red 80. Used for staining collagen and amyloid in histology.

25 g SR-P-25

Sudan III, C.I. 26100

Solvent Red 23, Sudan G, Fat Ponceau G, BSC certified stain. For lipid staining according to Lillie-Ashburn.

25 g S3-P-25

Sudan IV, C.I. 26105

Solvent Red 24, Scarlet Red, Biebrich Scarlet R, Fat Ponceau. For staining lipids according to Churukian's modification of Lillie-Ashburn's method.

25 g S4-P-25

Sudan Black B, C.I. 26150

Sudan Black, Solvent Black 3, Fat Black HB. For staining lipids and other hydrophobic materials.

5 g SB-P-5 10 g SB-P-10

Toluidine Blue O, C.I. 52040

Basic Blue 17, Tolonium Chloride, BSC certified stain. For staining mast cells, cartilage and Nissl bodies in the spinal cord.

25 g TB-P-25 100 g TB-P-100

Victoria Blue B, C.I. 44045

Basic Blue 26, Fat Blue B. For use in cytology and histology.

25 g VBB-P-25

Victoria Blue R, C.I. 42563

Basic Blue 8, Fat Blue 4R. For use in microbiology and histology.

25 g VBR-P-25

Water Blue, C.I. 42755

Acid Blue 22, BSC certified stain. For staining connective tissue according to Mallory and Gomori trichrome methods.

25 g WB-P-25

Wright's Stain

Colorants de Wright, BSC certified stain. Part of polychromatic Romanowsky stains commonly used in hematology.

25 g WS-P-25

Xylidine Ponceau, C.I. 16150

Ponceau 2R, Ponceau de Xylidine, BSC certified stain. For Masson trichrome staining.

25 g XP-P-25







Laboratory diagnostics

High quality disposable laboratory equipment includes products made of glass, plastic and metal, both sterilized and non-sterilized products, as well as single use and multiple use products. The products include VitroGnost slides for microscopy, cassettes for histopathology, sample transport containers and various other equipment and accessories.

- 64 VitroGnost slides for microscopy
- 67 Sample transport containers
- 68 Cassettes for histopathology
- 70 Pads and molds for histology
- 70 Equipment and accessories
- 71 Accessories for microtome
- 72 Jars and stands for staining microscopy sections
- 73 Accessories for storing microscope sections
- 74 Brushes
- 75 Laboratory equipment for urine sediment analysis
- 75 Other laboratory equipment

VitroGnost slides for microscopy

Super Grade

VitroGnost® Super Grade microscope slides

These slides are ideal for histology, pathology and cytology. Super Grade microscope slides of the highest quality and highest purity level ensure optimal conditions for microscopy and diagnostics.
Dimensions: 26x76x1 mm.

Cut edges, 45° corners	50 pcs	10117102
Cut edges, frosted both sides, 45° corners	50 pcs	10117107-1
Polished edges, 45° corners	50 pcs	10117101P
Polished edges, frosted both sides, 45° corners	50 pcs	10117107P
Polished edges, 45° corners	50 pcs	10127101P
Polished edges, frosted both sides, 45° corners	50 pcs	10127107P
Beveled edges, 45° corners	50 pcs	10127101A
Beveled edges, frosted both sides, 45° corners	50 pcs	10127107A

VitroGnost® Color Super Grade microscope

These slides are ideal for histology, pathology and cytology. Color Super Grade microscope slides of the highest quality and highest processing level ensure optimal conditions for microscopy and diagnostics. Colored label field of the slides enable easier sorting and marking.

Dimensions:	26x76x1	mm.
-------------	---------	-----

☐ Polished edges, WHITE, 45° corners	50 pcs	10128105PW
Polished edges, BLUE, 45° corners	50 pcs	10128105PB
Polished edges, YELLOW, 45° corners	50 pcs	10128105PY
Polished edges, GREEN, 45° corners	50 pcs	10128105PG
Polished edges, PINK, 45° corners	50 pcs	10128105PP
Polished edges, ORANGE, 45° corners	50 pcs	10128105PO
■ Polished edges, LILAC, 45° corners	50 pcs	10128105PL
Polished edges, PEACH, 45° corners	50 pcs	10128105PPe
Polished edges, PEACH, 45° corners Polished edges, GREY, 45° corners	50 pcs 50 pcs	10128105PPe 10128105PGr
	· ·	
Polished edges, GREY, 45° corners Polished edges, BROWN, 45°	50 pcs	10128105PGr



Economy Grade Standard Grade

VitroGnost® Economy Grade microscope slides

These slides are ideal for biochemistry and microbiology. Tissue paper interleaved. Economy Grade microscope slides are suitable for optical microscopy, routine procedures in biochemistry and microbiology laboratories. Dimensions: 26x76x1 mm.

Cut edges, 45° corners	50 pcs	7102
Cut edges, frosted one side, 90° corners	50 pcs	7105-1
Polished edges, 45° corners	50 pcs	7101
Polished edges, frosted one side, 45° edges	50 pcs	7105



Cover glass for use in microscopy

VitroGnost® cover glass - standard pack

0.13-0.17 mm thick microscope cover glass (Nr. 1.0). VitroGnost cover glasses of various dimensions used in optical microscopy.

18x18 mm	200 pcs	10211818C
20x20 mm	200 pcs	10212020C
22x22 mm	200 pcs	10212222C
24x24 mm	200 pcs	10212424C
24x32 mm	100 pcs	10212432C
24x40 mm	100 pcs	10212440C
24x50 mm	100 pcs	10212450C
24x60 mm	100 pcs	10212460C

VitroGnost® Standard Grade microscope slides

These slides may be used for hematology, pathology and cytology. Tissue paper interleaved. Standard Grade microscope slides are suitable for optical microscopy, routine procedures in hematology, cytology and pathology laboratories. Dimensions: 26x76x1 mm.

Cut edges, 45° corners	50 pcs	10217102
Cut edges, frosted both sides, 45° corners	50 pcs	10217107-1
Polished edges, 45° corners	50 pcs	10217101P
Polished edges, frosted both sides, 45° corners	50 pcs	10217107P
Polished edges, 45° corners	50 pcs	10227101P
Polished edges, frosted both sides, 45° corners	50 pcs	10227107P



VitroGnost slides for microscopy

Adhesive microscope slides

VitroGnost® Plus Ultra adhesive microscope slides

These slides are ideal for pathology, cytology, and special immunohistochemistry (IHC). VitroGnost Plus Ultra adhesive slide is a new generation slide with a denser and stronger positive charge on the surface to achieve even better adhesion of the sample. VitroGnost Plus Ultra adhesive slide can be used for many scientific purposes, especially with tissues difficult to fix, like breast tissue and samples of brain and skin. Dimensions: 26x76x1 mm.

72 pcs 103001A72

VitroGnost® PLL adhesive microscope slides

These slides are ideal for pathology, cytology, and special immunohistochemistry (IHC). Poly-L-lysine ensures covalent binding of the specimen to the microscope slide allowing better adhesion of the specimen onto slide. VitroGnost PLL adhesive microscope slides can be used with all samples regardless of the fixative used. Dimensions: 26x76x1 mm.

72 pcs 710972PLL

VitroGnost® SIL adhesive microscope slides

These slides are ideal for pathology, cytology, and special immunohistochemistry (IHC). Positively charged silanized VitroGnost adhesive microscope slides ensure better adhesion of negatively charged specimens onto slide. VitroGnost SIL adhesive microscope slides can be used with all samples regardless of the fixative used. Dimensions: 26x76x1 mm.

72 pcs 710972SIL

Slides for special use

ChlamyGnost DFA microscope slides

Microscope slides for microbiology. VitroGnost special microscope slides have circled marks for samples. The slides are of highest quality and purity level. DFA slides are used for direct fluorescent diagnostics of Chlamydia trachomatis. Dimensions: 26x76x1 mm.

50 pcs 103710950



Sample transport containers

Polypropylene containers for transporting samples

Container with cover for transporting histology samples and organs.

500 mL	1 pc	PK500
1000 mL	1 pc	PK1000
1200 mL	1 pc	PK1200
2500 mL	1 pc	PK2500
3750 mL	1 pc	PK3750
5000 mL	1 pc	PK5000
7000 mL	1 pc	PK7000
10000 mL	1 pc	PK10000

Polypropylene container with screw cap for histology

Plastic container with screw cap for sampling and transporting histology and cytology samples. Ideal for liquid specimen collection, transportation and storage in harsh conditions. Effectively prevents formalin leakage and evaporation. Complies with transportation requirements.

graded mark to	volume to the top	packaging	product code
7 mL	7 mL	1 pc	TP-7-7
20 mL	25 mL	1 pc	TP-20-20
40 mL	50 mL	1 pc	TP-40-40
60 mL	75 mL	1 pc	TP-60-60
90 mL	95 mL	1 pc	TP-90-90
120 mL	145 mL	1 pc	TP-120-120
160 mL	170 mL	1 pc	TP-160-160
250 mL	275 mL	1 pc	TP-250-150
1000 mL	975 mL	1 pc	TP-1000-750



Cassettes for histopathology

Cassettes without lid - standard

Standard cassettes without lids made of polyoxymethylene with writing area for easy sample marking, organizing and identifying. Available in 5 different colors.

color	material	packaging	product code
White	P.O.M.	500 pcs	103W
Yellow	P.O.M.	500 pcs	103Y
Green	P.O.M.	500 pcs	103G
Pink	P.O.M.	500 pcs	103P
Blue	P.O.M.	500 pcs	103B



Biopsy cassettes with lidBiopsy cassettes made of polyoxymethylene with writing area for easy sample marking, organizing and identifying. Available in 12 different colors.

color	material	packaging	product code
White	P.O.M.	500 pcs	101W
Blue	P.O.M.	500 pcs	101B
Yellow	P.O.M.	500 pcs	101Y
Green	P.O.M.	500 pcs	101G
Pink	P.O.M.	500 pcs	101P
Orange	P.O.M.	500 pcs	1010
Lilac	P.O.M.	500 pcs	101L
Peach	P.O.M.	500 pcs	101Pe
Grey	P.O.M.	500 pcs	101Gr
Brown	P.O.M.	500 pcs	101Br
Aqua	P.O.M.	500 pcs	101Aq
Beige	P.O.M.	500 pcs	101Be



Lid for cassettes

Cassette embedding stainless steel lid, standard.


material	packaging	product code
INOX	1 pc	51050103



Histology cassettes with lidHistology cassettes with lid made of polyoxymethylene with writing area for easy sample marking, organizing and identifying. Available in 12 different colors.

TYPE 102 - closing at the front side of the lid

color	material	packaging	product code
White	P.O.M.	250 pcs	102W
Blue	P.O.M.	250 pcs	102B
Yellow	P.O.M.	250 pcs	102Y
Green	P.O.M.	250 pcs	102G
Pink	P.O.M.	250 pcs	102P
Orange	P.O.M.	250 pcs	1020
Lilac	P.O.M.	250 pcs	102L
Peach	P.O.M.	250 pcs	102Pe
Grey	P.O.M.	250 pcs	102Gr
Brown	P.O.M.	250 pcs	102Br
Aqua	P.O.M.	250 pcs	102Aq
Beige	P.O.M.	250 pcs	102Be



color	material	packaging	product code
White	P.O.M.	500 pcs	107W
Blue	P.O.M.	500 pcs	107B
Yellow	P.O.M.	500 pcs	107Y
Green	P.O.M.	500 pcs	107G
Pink	P.O.M.	500 pcs	107P
Orange	P.O.M.	500 pcs	1070
Lilac	P.O.M.	500 pcs	107L
Peach	P.O.M.	500 pcs	107Pe
Grey	P.O.M.	500 pcs	107Gr
Brown	P.O.M.	500 pcs	107Br
Aqua	P.O.M.	500 pcs	107Aq
Beige	P.O.M.	500 pcs	107Be

Cassettes with lid for small biopsy

Small biopsy cassettes with lid made of polyoxymethylene with writing area for easy sample marking, organizing and identifying.

color	material	packaging	product code
White	P.O.M.	250 pcs	104W
Blue	P.O.M.	250 pcs	104B
Yellow	P.O.M.	250 pcs	104Y
Green	P.O.M.	250 pcs	104G
Pink	P.O.M.	250 pcs	104P
Orange	P.O.M.	250 pcs	1040
Lilac	P.O.M.	250 pcs	104L
Peach	P.O.M.	250 pcs	104Pe
Grey	P.O.M.	250 pcs	104Gr
Brown	P.O.M.	250 pcs	104Br
Aqua	P.O.M.	250 pcs	104Aq
Beige	P.O.M.	250 pcs	104Be







Pads and molds for histology

Metal histology mold

dimension	material	packaging	product code
7x7x6 mm	INOX	1 pc	51050707
15x15x6 mm	INOX	1 pc	51051515
24x24x6 mm	INOX	1 pc	51052424
30x24x6 mm	INOX	1 pc	51053024
37x24x6 mm	INOX	1 pc	51053724
65x45x15 mm	INOX	1 рс	BO07-7010

Biopsy pads for cassettes, blue Solvent resistant biopsy pads for small sample processing.

1000 pcs SP-1000



Equipment and accessories

Polypropylene dissecting boardsBoards for pathohistological tissue dissection made of polypropylene.

Polypropylene, white, dimensions 60x40x1.5 cm	1 pc	BC001
Polypropylene, dark blue, dimensions 39x29.5x2.5 cm	1 pc	BC032
Polypropylene, white, dimensions 40x30x1.5 cm	1 pc	BC061CS



Accessories for microtome

Microtome blades for histology, high profile

Bio/Cut microtome blades - high profile

Bio/Cut H35C microtome blades for cryotome, high profile (wide). Ideal for frozen sections. Suitable for both soft and hard tissues.

50 pcs Bio/Cut H35C

Histoblades Duraedge histology blades - high profile

HistoBlades Duraedge blades with Teflon coating - high profile.

50 pcs 7310

Microtome blades for histology, low profile

Bio/Cut microtome blades - low profile

Bio/Cut microtome blades manufactured in order to provide most precise sectioning.

Bio/Cut 22A microtome blades for ultra-thin sections, kidneys and lymph nodes, narrow blade	50 pcs	Bio/Cut 22A
Bio/Cut 35A microtome blades for soft and hard tissues, low profile	50 pcs	Bio/Cut 35A
Bio/Cut 35B microtome blades for soft tissues, low profile	50 pcs	Bio/Cut 35B
Bio/Cut 35N microtome blades for thin sections, low profile	50 pcs	Bio/Cut 35N
Bio/Cut 35R microtome blades for hard tissues, low profile	50 pcs	Bio/Cut 35R
Bio/Cut 35S microtome blades for hard tissues, low profile	50 pcs	Bio/Cut 35S

Histoblades Duraedge histology blades - low profile

HistoBlades Duraedge blades with Teflon coating - low profile.

50 pcs 7223

Histoblades Encore histology blades - low profile

Extremely sharp new generation Teflon coated blades - low profile.

50 pcs 7380





Accessories for microtome

Additional products for microtome

Microtome oil

Optimal viscosity oil for microtome maintenance.

Microtome oil 50 mL	1 pc	MO-50
Microtome oil 100 mL	1 pc	MO-100



Jars and stands for staining microscopy sections

Glass staining jarsGlass staining jars with covers for staining according to Coplin, Hellendahl and Schifferdecker.

Glass staining jars for 5 slides with cover, Coplin	1 pc	10130505
Glass staining jars for 9 slides with cover, Hellendahl	1 pc	10130609
Glass staining jars for 10 slides with cover, Schifferdecker	1 pc	10130510

Glass jars and metal racks for staining 30 - 60 samples

Glass jars (transparent or brown) with covers for staining samples and corresponding racks.

Glass staining jars for 30 slides with cover (with rack, product code: 50101630)	1 pc	10130530C
Glass staining jars for 30 slides with cover, brown (with rack, product code: 50101630)	1 pc	10130530B
Glass staining jars for 60 slides with cover (with rack, product code: 50101660)	1 pc	10130560
Rack for staining 30 slides, inox (for glass jars, product code: 10130530C and B)	1 pc	50101630
Rack for staining 60 slides, inox (for glass jars, product code: 10130560)	1 рс	50101660



Accessories for storing microscope sections

Microscope slide mailers

Microscope slide mailers made of polypropylene. Ideal for microscope slide transport and storage. Made of thick, high-strength material.

Microscope slide mailer for 1 slide, polypropylene	100 pcs	31011001MA
Microscope slide mailer for 2 slides, polypropylene	100 pcs	31011002MA
Microscope slide mailer for 2 slides, polypropylene, blue, slides in vertical position	100 pcs	31011102BB
Microscope slide mailer for 2 slides, polypropylene, colorless, slides in vertical position	100 pcs	31011102BN
Microscope slide mailer for 5 slides, polypropylene, colorless, slides in vertical position	100 pcs	31011005MA

Microscope slide plastic box

Boxes for microscope slide storage. Compact and long-lasting boxes are the best choice for slide protection in laboratory. Enumerated stands make slide organization easier and simpler. Made of ABS material.

Microscope slide box for 12 slides, ABS, orange	1 pc	31012012AO
Microscope slide box for 25 slides, ABS, red, blue or green	1 pc	31012025AG
Microscope slide box for 50 slides, ABS, grey	1 pc	31012050AG
Microscope slide box for 100 slides, ABS, grey or white	1 pc	31012100AG

Cardboard microscope slides tray

Microscope slide tray made of cardboard used for holding and storing histology slides.

Cardboard microscope slide tray for 20 slides, without cover	10 pcs	60001021
Cardboard microscope slide tray for 20 slides, with cover and partitions	5 pcs	60001020
Cardboard microscope slide tray for 24 slides, with cover and without partitions	5 pcs	60001024

Plastic microscope slides trayMicroscope slide tray made of plastic used for holding and storing histology slides.

Plastic microscope slide tray for 20 slides, without cover, white	1 pc	31013020W
Plastic microscope slide tray for 20 slides, without cover, green	1 pc	31013020G
Plastic microscope slide tray for 20 slides, without cover, pink	1 pc	31013020P
Plastic microscope slide tray for 20 slides, without cover, blue	1 рс	31013020B



Brushes

Cervical sampling tools

Cervical Brush

Disposable brushes for cytological sample collection.

Cervical Brush for cervical samples collection

50 pcs

92014000

Cervical Brush Combi

Disposable brushes for ectocervical and endocervical samples collection with a snap-off head.

Cervical Brush Combi for cervical samples collection

50 pcs

92013000

Cervical Brush S

Disposable brushes for vaginal, ectocervical and endocervical samples collection.

Cervical Brush S for cervical samples collection

50 pcs

92011000

Cervical spatula

Disposable Ayre spatulas for ectocervical and endocervical sampling.

Cervical spatula 100 pcs 93021000

Cito Brush

Disposable brushes for endocervical samples collection.

Cito Brush for cervical samples collection 100 pcs 92010000

Cito Brush for cervical samples 100 pcs 92010000EO collection, sterile



Laboratory equipment for urine sediment analysis

Urine tubes and pipettes for urine sediment analysis

Tubes with retentive bottom for urine sedimentation and pipette for decanting of sediment.

UriGnost tube, 15 mL, with screw 200 pcs 84020-2501 cap, retentive bottom, vol. 0.2 mL, graduated at 2.5/5/10/12 mL

Pipette tips 200 μ L, Eppendorf/ 500 pcs 3601-500 Universal type, yellow

Urine plates for urine sediment analysis

Disposable plates used for microscopical analysis and counting elements of urine sediment.

UriGnost 10 plate, 50 pcs for 500 urine sediment analyses	50 pcs	UG10P-50
UriGnost 10 plate, 100 pcs for 1000 urine sediment analyses	100 pcs	UG10P-100
* part of UriGnost S and UriGnost SM kits, for more information, see		

Other laboratory equipment

Postmortem suture

Postmortem polyester suture.

Postmortem suture, polyester, white	spool length 450 m	KZS BA028
Postmortem suture, polyester 1 mm, processed with wax, beige	spool length 335 m	KZS BA024
Postmortem suture, polyester 1 mm, processed with wax, white	spool length 335 m	KZS BA041

Histology filter paper - sheets

For general laboratory use.

pages 50 and 57

Filter papers for histology - sheets 1 pc 22/N size 50x50 cm

 $^{^{\}star}$ part of UriGnost S and UriGnost SM kits, for more information, see pages 50 and 57

Contact information

BioGnost Ltd., Medjugorska 59, Zagreb 10040 Croatia (Europe)

WORKING HOURS Monday through Friday 8:00 to 16:00 (CET)

WORLDWIDE DISTRIBUTION: Contact us to find your local distributor. BioGnost is currently shipping to over 70 countries.



Ordering information

e-mail: sales@biognost.hr website: www.biognost.com phone: +385 1 2409 997 fax: +385 1 2404 039

To help us make sure that your order is processed and delivered correctly, please have the following information available:

- Name of your organization
- Post code
- Contact name, phone number and e-mail address
- · Delivery and Invoice address
- · Catalog number
- Quantity

Can't find the product you require? BioGnost may offer what you're looking for as a non-stock item or a custom offer.



Technical support

Technical questions can be answered by emailing our sales representatives at sandra.ban@biognost.hr, pavle.loncar@biognost.hr and viktorija.liber@biqnost.hr.

They will provide you with any MSDS or product information required, also available on our website at www.biognost.com.

Frequently Asked Questions (FAQ) section is available on our website.



Shipping information

Shipping and handling charges are prepaid and added to the invoice. If requested, we can use your account number or carrier of your choice, and also ship on your third-party freight accounts (Incoterms 2010). Charges vary according to the destination, weight and content, and are available upon request.



Our employees are trained to classify, package, mark and label products identified as dangerous goods according to ADR, IMDG and IATA to ensure safe transport

If you require more information about shipping, please contact our Logistics Department at sinisa.carevic@biognost.hr.

A full transcript of the current Terms and Conditions is available on BioGnost website at www.biognost.com.

T	
In	$-\mathbf{x}$

Troples		BioMount C	19
Index		BioMount DPX	17
		BioMount DPX High	17
		BioMount DPX Low	17
4% Formaldehyde	10	BioMount DPX New	17
4% NB Formaldehyde	10	BioMount High	18
10% Formaldehyde	11	BioMount M	18
10% Formalin, alcoholic solution	52	BioMount New	17
10% NB Formaldehyde	11	BioMount New Low	17
		Biopsy cassettes with lid	68
Α.		Biopsy pads for cassettes, blue	70
A		BioSchiff reagent	52
Acetic acid, 0.5% solution	52	BioWax 52/54	22
Acid alcohol	28	BioWax 56/58	22
Acid alcohols	52	BioWax Blue	22
Acid solution	52	BioWax Micro	22
Acridine Orange, C.I. 46005	58	BioWax Plus 56/58	22
Acridine Orange solution	52	Bismarck Brown Y, C.I. 21000	58
A.F.O.G. kit	43	Bluing reagent	28
A.F.O.G. reagent	52	Borax, solution	52
Alcian Blue 8 GX, C.I. 74240	58	Bouin's solution	12 58
Alcian Blue - P.A.S. kit	43	Brilliant Cresyl Blue, C.I. 51010	
Alcian Blue pH 1.0 kit	43	Brilliant Cresyl Blue, solution Brilliant Green Certified, C.I. 42040	35 58
Alcian Blue pH 2.5 kit	43	·	
Alcian Blue solution 1%	52	Buffer solution, pH 6.4	34
Alcian Blue solution pH 1.0	52	Buffer solution, pH 6.8	34
Alcian Blue solution pH 2.5	52	Buffer solution, pH 7.2 Buffer tablets pH 6.8	33 34
Alcian Green	58		33
Alcian Yellow, C.I. 12840	58	Buffer tablets pH 7.2	33
Alcian Yellow Toluidine Blue kit	43		
Alcoholic Eosin Y solutions	29		
Alkaline alcohol, solution	52		
Amido Black 10B, C.I. 20470	58	Canada Balsam	19
Ammonium iron sulfate, solution	52	Carbol Fuchsin Gram 10x concentrated	37
Aniline, alcoholic solution	52	Cardboard microscope slides tray	73
Aniline Blue, C.I. 42780	58	Carmine, C.I. 75470	58
Aniline Blue reagent	52	Carminic acid, C.I. 75470	58
Aniline Blue, solution	52	Cassettes with lid for small biopsy	69
Auramine O, C.I. 41000	58	Cassettes without lid - standard	68
Azan reagent	52	Cedarwood oil	21
Azan Trichrome kit	43	Celestine Blue, C.I. 51050	59
Azocarmine G, C.I. 50085	58	Cervical Brush	74
Azocarmine, solution	52	Cervical Brush Combi	74
Azophloxine, C.I. 18050	58	Cervical Brush S	74
Azure A, C.I. 52005	58	Cervical spatula	74
Azure B, C.I. 52010	58	ChlamyGnost DFA microscope slides	66
Azure C, C.I. 52002	58	Chromotrope 2R / Aniline Blue reagent	52
Azure II	58	Chromotrope 2R, C.I. 16570	59
Azure II Eosinate	58	Chromotrope 2R / Fast Green F.C.F. reagent	53
		Cito Brush	74
D		CitoFix	12
В		CitoSpray	12
Basic Activation buffer	52	CitoSpray solution	12
Benedict's reagent	52	Clean Lens, solution 1	21
Biebrich Scarlet-Acid Fuchsin reagent	52	Clean Lens, solution 2	21
Biebrich Scarlet, C.I. 26905	58	Clean Lens, solution 3	21
BioClear	15	Colloidal Iron kit	44
BioClear New	15	Colloidal Iron, stock solution	53
Bio/Cut microtome blades - high profile	71	Congo Red, C.I. 22120	59
Bio/Cut microtome blades - low profile	71	Congo Red Highman kit	44
Bio-Diff 1 reagent	32	Congo Red Puchtler kit	44
Bio-Diff 1 RTU reagent	33	Congo Red Puchtler reagent	53
Bio-Diff 2 reagent	32	Congo Red reagent	53
Bio-Diff 2 RTU reagent	33	Cresyl Violet Acetate Certified	59
Bio-Diff 3 reagent	32	Cresyl Violet, solution	53
Bio-Diff 3 RTU reagent	33	CryoFix Gel	23
Bio-Diff kit	32	Cryo Spray	23
Bio-Diff RTU kit	33, 44	Crystal Violet, C.I. 42555	59
BioFix	11		
BioFix C	11	D	
BioFix GL	11	D	
BioFix HD	11	DeHematin	12
BioForm S	11	DeMold solution	23
BioGram 4 kit	36	DeMold Spray	23
BioGram ECO kit	36	Differentiation reagents based on acetic acid	53
BioGram Histo kit	44	Differentiation reagents based on hydrochloric acid	53
BioMount	18	Duoline marker pen	9
BioMount Aqua	19	ı	
the state of the s			

E		Hematoxylin H	27
	24	Hematoxylin HP, Pap 1A	30
EA 31 reagent, Pap 3A	31	Hematoxylin HP, Pap 1B	30
EA 50 reagent, Pap 3B	31	Hematoxylin Instant kit	28
EA 65 reagent, Pap 3C	31	Hematoxylin M	27
EA 65 reagent, Pap 3D	31	Hematoxylin ML	27
Ehrlich aldehyde reagent	53	Hematoxylin P.T.A.	28
Elastica-Van Gieson kit	44	Hematoxylin P.T.A. kit	46
Eosin and nigrosine vital kit	45	Hematoxylin, Verhoeff A	54
Eosin B, C.I. 45400	59	Hematoxylin, Weigert A	54
Eosin Contrast	29		8, 46
Eosin Contrast PLUS	29	HemoGnost Perls kit	46
Eosin-nigrosin vital	53	Histanol 50	14
Eosin Y aqueous solutions	29	Histanol 70	14
Eosin Y, C.I. 45380	59	Histanol 80	14
Eosin Y instant	29	Histanol 95	14
Eosin Y, isotonic solution	54	Histanol 96	14
Eriochrome Cyanine R	59	Histanol 100	14
		Histanol EP 95	14
E. Carriera		Histanol EP 100	14
F		Histanol G	14
Fast Green F.C.F., C.I. 42053	59	Histanol IP	14
Fast Green F.C.F. contrast reagent	54	Histanol M	14
Fast Green F.C.F. reagent	54	Histoblades Duraedge histology blades - high profile	71
Ferri reagens, Verhoeff B	54	Histoblades Duraedge histology blades - low profile	71
Ferri reagent, Weigert B	54	Histoblades Encore histology blades - low profile	71
Feulgen kit	45	Histology cassettes with lid	69
Field kit	35, 4 5	Histology filter paper - sheets	75
Field, solution A	54	Hydroquinone, solution	54
Field, solution B	54		
Formaldehyde 37% for use in histology	11		
Fouchet-Van Gieson kit	45	I	
Fuchsin Acid, C.I. 42685	59	_	20
	54	Immersion oil	20
Fuchsin Acid Van Ciasan reagent		Immersion oil, tropical grade	21
Fuchsin Acid Van Gieson reagent	54 59	Immersion oil type 37	21
Fuchsin Basic, C.I. 42510		Immersion oil type A	20
Fuchsin Basic reagent	54	Immersion oil type C	21
Fuchsin Basic - Schiff, C.I. 42500	59	Immersion oil type FF	21
Fuchsin Basic, solution	54	ImmunoForm	11
		Iodine solution, Verhoeff C	54
		Iron chloride, solution	54
G		Iron nitrate, solution	54
Gabbet reagent	39		
Gelatin, gel in transport jar	54	The second secon	
Gentian Violet, C.I. 42555	59	J	
Giemsa	59	Janus Green B, C.I. 11050	59
Giemsa HP kit	45	Jenner Stain	59
Giemsa solution	34		
Glass jars and metal racks for staining 30 - 60 samples	72		
Glass staining jars	72	K	
Glutaraldehyde, 4% buffered solution	54		
Gold chloride solutions	54	Kit for rapid HE staining of frozen and paraffin-embedded sections	
Gomori Trichrome kit	45	Kit for rapid Pap staining	30
Gram Crystal Violet 1% solution	36	Kovac indole reagent	54
Gram Crystal Violet, phenol-free reagent	36		
Gram Decolorizer solution 1	37	The second secon	
Gram Decolorizer solution 2	36	L	
Gram Decolorizer solution 3	37	Lactophenol Blue solution	41
Gram Lugol solution	37	Lactophenol solution	41
Gram Lugol solution, stabilized	36	Leishman's solution	35
Gram Safranin 5x concentrated	37	Leishman Stain	59
Gram Safranin solution	36	Lid for cassettes	68
Gram sodium hydrogencarbonate, solution	36	Light Green S.F., C.I. 42095	59
Grimelius kit	46	Light Green S.F. reagent	54
Grocott kit	46	Liquid Blocker marker pen	9
GIOCOTT INT		Lithium carbonate, Luxol	54
		Lithium carbonate, Euxor Lithium carbonate, solution	54
H		Lugol's C solution	37
		Lugol's solution, acid	37
Hayem's solution	35	Lugol's solution, Acid Lugol's solution, M.I.F.	55
H.B.F.P. kit	46	Lugol's solution, M.I.F. Lugol's solution (Schiller's test)	37
HCL Pap reagent	28	Luxol Fast Blue kit	47
Hematoxylin CAR	28	Luxol Fast Blue MBSN, C.I. 74180	59
Hematoxylin, C.I. 75290	59		59 55
Hematoxylin COL	28	Luxol Fast Blue, solution	22
Hematoxylin G1	27, 30		
Hematoxylin G2	27, 30		
Hematoxylin G3	27, 30		

N A		Phosphomolybdic acid, 5% solution	56
M		Phosphotungstic acid, 1% solution	56
Malachite Green, C.I. 42000	60	Phosphotungstic acid, 5% solution	56
Mallory Trichrome kit	47	Picric acid in acetone, solution	56
Martius Scarlet Blue (MSB) kit	47	Plastic microscope slides tray	73
Martius Yellow C.I. 10315	60	Polypropylene containers for transporting samples	67
Martius Yellow, solution	55	Polypropylene container with screw cap for histology	67
Masson Fontana kit	47	Polypropylene dissecting boards	70
Masson-Goldner Trichrome kit	47	Ponceau S, C.I. 27195	60
Masson Trichrome kit	48	Ponceau S, solution	56
Mayer's mucicarmine	55	Postmortem suture	75
May-Gruenwald	60	Potassium hexacyanoferrate, Colloidal Iron solution	56
May-Gruenwald solution	34	Potassium hexacyanoferrate, solution	56
Metal histology mold	70	Potassium hydroxide, solution	56
Metanil Yellow, C.I. 13065	60	Potassium hydroxide, solution for fungi	41
Metanil Yellow, solution	55	Potassium permanganate	60
Methenamine, solution	55	Potassium permanganate, solutions	56
Methyl Blue Eosin, solution	55	Propylene glycol	56
Methylene Blue 10X concentrated	38	Propylene glycol, 85%	56
Methylene Blue, C.I. 52015	60	P.T.A P.M.A. reagent	56
Methylene Blue Loeffler's reagent	38	Pyronine B, C.I. 45010	60
Methylene Green, C.I. 52020	60	,	
Methyl Green, C.I. 42590	60		
Methyl Violet 2B, C.I. 42535	60	R	
Methyl Violet 10B, C.I. 42555	60		
M.F. reagent	55	Reagent for differentiation in Verhoeff kit	56
Micro Clean	22	Reducing buffer, solution	56
Microscope slide mailers	73	Resorcine Fuchsin reagent	56
Microscope slide plastic box	73	Resorcinol, C.I. 76505	60
Microtome oil	72	Reticulin Contrast kit	49
M.I.F. kit	47	Reticulin kit	49
Movat kit	48	Rhodamine B, C.I. 45170	60
Mucicarmine kit	48	Rhodanine	61
Myco-Mount Blue	41	Rhodanine kit	49
Myco-Mount Red	41	Rhodanine reagent	56
,		Rose Bengal, C.I. 45440	61
NI			
N		S	
NB Formaldehyde concentrated 5x	11		
Neutral Red, C.I. 50040	60	Saccomanno fixative	12
Neutral Red reagent	55	Safranin O, C.I. 50240	61
Nigrosine, isotonic solution	55	Scott's solution	28
Nigrosine WS, C.I. 50420	60	Shorr's solution	30
Nuclear Fast Red, C.I. 60760	60	Silver nitrate solutions	56
Nuclear Fast Red (Kernechtrot) reagent	55	Sirius Red, C.I. 35780	61
(Sodium acetate, solution	56
		Sodium hydroxide, solution	56
lacksquare		Sodium metabisulfite solutions	56
		Sodium tetraborate, solution	57
OG-6 reagent, Pap 2A	30	Sodium thiosulfate solution	57
OG-EA31 Pap reagent	31	Sperm-Diff RTU kit	49
OG-EA50 Pap reagent	31	Sudan Black B, C.I. 26150	61
Oil Red O, C.I. 26125	60	Sudan Black B kit	50
Oil Red O kit	48	Sudan Black B lipid kit	50
Oil Red O reagent	55	Sudan Black B reagent	57
Orange G, 1% solution	55	Sudan Black B, solution	57
Orange G / Aniline Blue reagent	55	Sudan III, C.I. 26100	61
Orange G, C.I. 16230	60	Sudan IV, C.I. 26105	61
Orange II, C.I. 15510	60	Sulfuric acid solutions	57
Orange II reagent, Pap 2B	30	Sulphosalicylic acid, 20% solution	57
Orcein, C.I. (ed.1) 1242	60		
Orcein kit	48	_	
Orcein reagent	55		
OsteoFast 1	13	TB Armand reagent	39
OsteoFast 2	13	TB Auramine O reagent	40
OsteoSens	13	TB Auramine O reagent TB Auramine-Rhodamine reagent	40
Oxalic acid solutions	55		
		TB Carbol Fuchsin reagent TB Decolorizer	38, 39 38, 39
D		TB Decolorizer TB Decolorizer Fluorescent	30, 39 40
P			40 38
Paraldehyde Fuchsin kit	48	TB-Fuchsin reagent	
Paraldehyde Fuchsin reagent	55	TB Malachite Green reagent	39
P.A.S. kit	49	TB Permanganate reagent	40
P.A.S.M. / Jones kit	49	TB-Stain Auramine O kit	40
		TB-Stain Cold kit	39
Periodic acid, solution	55 60	TB-Stain ECO kit	38
Phosphate buffered saline (PBS)	60 55	TB-Stain Fluorescent kit	40
Phosphomolyhdic acid 1% solution	55 56	TB-Stain Histo kit	50
Phosphomolybdic acid, 1% solution	50		

TB-Stain Hot kit	38
TB-Stain Quick kit	39
Tincture of iodine, 1%	57
Tissue marking dyes	8
Tissue marking dyes, 5 colour set	8
Tissue marking dyes, 7 colour set	8
Tissue Soft	23
Toluidine Blue O, C.I. 52040 Toluidine Blue, solution	61 57
Trichloroacetic acid, solution	57 57
Tuerk's solution	35
Tachka solution	
H.	
O .	
UriGnost S kit	50
UriGnost SM kit	50
UriGnost SM reagent	57
UriGnost S reagent	57
Urine plates for urine sediment analysis	75 75
Urine tubes and pipettes for urine sediment analysis	/5
V	
Van Gieson Trichrome kit	50
Verhoeff kit	51
Victoria Blue B, C.I. 44045	61
Victoria Blue R, C.I. 42563	61
VitroGnost® Color Super Grade microscope slides	64
VitroGnost® cover glass - standard pack	65
VitroGnost® Economy Grade microscope slides	65
VitroGnost® PLL adhesive microscope slides	66
VitroGnost® Plus Ultra adhesive microscope slides	66
VitroGnost® SIL adhesive microscope slides	66
VitroGnost® Standard Grade microscope slides	65
VitroGnost® Super Grade microscope slides	64
Von Kossa kit	51
W	
Warthin Starry kit	51
Water Blue, C.I. 42755	61
Weigert-Van Gieson kit	51 35
Wright-Giemsa solution Wright-Giemsa, solution for reptiles	35 35
Wright's solution	35 35
Wright's Stain	61
9	01
Y	
VIII D CLICIFO	-
Xylidine Ponceau, C.I. 16150	61

NOTES	

NOTES	

