

BIOGNOST®
MEDICAL PRODUCTS


**MICROSCOPY AND
LABORATORY DIAGNOSTICS**



Our global presence

“ We manufacture products for use in histopathology, hematology, cytology, and microbiology with a wide national and international distribution network present in over 80 countries.

BioGnost offers high quality products at competitive prices and has one of the widest ranges of medical products in South-Eastern Europe. Our experience and flexibility towards the client makes us a sought-after business partner, which has been recognized on the global market.



Our entrepreneurship started in 1990 with the founding of the family company. Accelerated business growth began in 1998 with production of blood typing serums followed by the Croatian Ministry of Health granting BioGnost the License for the manufacture of medical products. We are proud of one of our biggest successes - relocating to our new commercial and manufacturing building in 2012. The site enabled an enhanced work environment, including a carefully devised warehouse structure, segmented production facilities with increased manufacturing capacity, and additional R&D capabilities.

“ We continue to provide high-quality products and supportive customer service.

Strategic priorities

Our mission is to create high-quality products and provide excellent services, as well as introduce new or expanded product lines and sales ranges in accordance with market dynamics and demands.

The key to our success is the ever-expanding network of business partnerships, as well as recognizable social responsibility. We plan to expand our market, increase investments in research and development, and establish our company as a renowned manufacturer in the fields of microbial diagnostics and histopathology.

ISO 9001, ISO 13485, ISO 14001

BioGnost's entire business is in accordance with international ISO 9001, ISO 13485 and ISO 14001 standards. The company implemented ISO standards more than 15 years ago, and is the first Croatian company to implement the ISO 13485 quality system for medical devices. Having fulfilled the requirements of these standards, the company is completely in accordance with national legal regulation and national legal regulation and the Regulation (EU) 2017/745 on medical devices as well as the Regulation (EU) 2017/746 on in vitro diagnostic medical devices. All of our IVD products (more than a thousand) have been listed in EUDAMED database.



CE, IVDR

BioGnost's products bear the CE mark, a certificate that confirms that the products are in accordance with the European Union's standards. They are declared as *in vitro* diagnostic medical devices (IVD) classified as "Class A" products (according to the Regulation (EU) 2017/746). No control of a Notified Body is required for this class; entire company business is in accordance with applicable regulations and standards.

OEM manufacture

Besides products from our own sales range, we also offer OEM manufacture for your company. It includes a wide range of services, such as label design, product packaging, instructions for use and certificates of analysis, and products manufactured according to your specifications.

Contents



FIXATION AND PROCESSING 6

Marking systems 6

Formalin fixatives 8

Other fixatives 10

Decalcifying solutions 12

Dehydrating and rehydrating agents 13

Clearing and deparaffinization agents 14

Infiltration, embedding and demolding media 15

MOUNTING AND MICROSCOPY 17

Mounting media 17

Immersion oils 21

Cleaning solutions for microscope lens and slides 22

STANDARD HISTOLOGY STAINING REAGENTS 23

Hematoxylin reagents 23

Differentiation and bluing reagents for histology and cytology 26

Eosin reagents 27

CYTOLOGY AND HEMATOLOGY STAINING REAGENTS 28

Papanicolaou (Pap) reagents 28

Rapid and standard hematology reagents and kit 30

LEUKOGNOST Cytochemical staining kits and ancillary reagents 34

BACTERIOLOGY AND MYCOLOGY STAINING REAGENTS 36

BioGram kits and reagents 36

TB-Stain kits and reagents 38

Solutions for staining and analysis of fungi 41

SPECIAL STAINING KITS FOR HISTOLOGY AND CYTOLOGY 42

ADDITIONAL STAINING REAGENTS AND SOLUTIONS 53

POWDER DYES AND STAINS 60

ACCESSORIES 64

Legend:

● DG cargo in transport ○ DG cargo only in air transport ● temperature controlled cargo (transport regime 2-8°C)



Formalin fixatives

Fixation is one of the most important steps in tissue processing, preventing tissue degradation and decomposition. Preserving tissue properties is important for detection of morphological and pathological changes in tissue samples.

4% NB Formaldehyde

10% neutral buffered and stabilized formalin solution, pH 7.0.

1000 mL	FNB4-1L
5000 mL	FNB4-5L
10000 mL	FNB4-10L
20000 mL	FNB4-20L



4% Formaldehyde

Stabilized 10% formalin solution.

1000 mL	F4-1L
5000 mL	F4-5L
10000 mL	F4-10L
20000 mL	F4-20L





FormSeal kit

10% neutral buffered and stabilized formalin solution in a closed-circuit system, for fixation and transport of small tissue samples and material from biopsies.

25x35 mL	FS-K-35
25x20 mL	FS-K-20



NEW

4% NB Formaldehyde in transport containers

10% neutral buffered and stabilized formalin solution in transport containers for fixation and transport of small tissue samples and material from biopsies.

25x35 mL	F4-60-35
25x20 mL	F4-40-20



10% NB Formaldehyde

25% neutral buffered and stabilized formalin solution, pH 7.0.

1000 mL	FNB10-1L
5000 mL	FNB10-5L
10000 mL	FNB10-10L
20000 mL	FNB10-20L



ImmunoForm

Stabilized 10% formalin solution produced from paraformaldehyde without methyl alcohol. For use in histology and immunohistochemistry.

1000 mL	IMF-1L
5000 mL	IMF-5L
10000 mL	IMF-10L



10% Formaldehyde

Stabilized 25% formalin solution.

1000 mL	F10-1L
5000 mL	F10-5L
10000 mL	F10-10L
20000 mL	F10-20L



Formaldehyde 37% for use in histology

100% formalin solution stable at low temperatures.

1000 mL	F37H-1L
5000 mL	F37H-5L
10000 mL	F37H-10L



NB Formaldehyde concentrated 5x

Concentrated neutral buffered formalin for preparation of 10% buffered formalin (synonym: 4% buffered formaldehyde) solution, pH 7.0.

For preparation of 5 L of 4% buffered formaldehyde	1000 mL	FK5X-1L
--	---------	---------

For preparation of 25 L of 4% buffered formaldehyde	5000 mL	FK5X-5L
---	---------	---------

For preparation of 50 L of 4% buffered formaldehyde	10000 mL	FK5X-10L
---	----------	----------



DeHematin

Modified Kardasewitsch solution for depigmentation of tissues fixed in non-buffered formalin (after acid hematin occurs).

1000 mL	KO-OT-1L
5000 mL	KO-OT-5L





Other fixatives

BioFix

Methyl alcohol-based fixative for cryostat sections.

500 mL	BF-500
1000 mL	BF-1L



BioFix C

Modified Carnoy's fixative based on ethyl alcohol, chloroform and acetic acid.

500 mL	BFC-500
1000 mL	BFC-1L
5000 mL	BFC-5L



BioFix GL

Glyoxal-based, formalin-free fixative. A less toxic/carcinogenic alternative to formalin fixatives. Suitable for all tissue types. For use in histology, immunohistochemistry, special stains and genetic analyses. Stable for 12 months at 2-8 °C or one month at 15-25 °C.

500 mL	BFGL-500
1000 mL	BFGL-1L
2500 mL	BFGL-2.5L
5000 mL	BFGL-5L



BioFix HD

Modified Hartmann/Davidson's fixative that enables rapid fixation and better visualization of lymph nodes. The results provide clearer nuclear details and minimize formalin pigment occurrence.

500 mL	BFHD-500
1000 mL	BFHD-1L
5000 mL	BFHD-5L



BioForm S

Modified Schaeffer's buffered fixative solution of formaldehyde in methanol.

1000 mL	BIOS-OT-1L
5000 mL	BIOS-OT-5L



ALSO AVAILABLE IN TRANSPORT CONTAINERS

BioFix GL in transport containers

Glyoxal-based, formalin-free fixative for fixation and transport of small tissue samples and material from biopsies.

25x35 mL	BFGL-60-35
25x20 mL	BFGL-40-20



Other fixatives

Bouin's solution

Aqueous solution of picric acid, formaldehyde and acetic acid for tissue fixation and better visualization of lymph nodes.

100 mL	BOU-OT-100
1000 mL	BOU-OT-1L
5000 mL	BOU-OT-5L
10000 mL	BOU-OT-10L



CitoFix

Alcoholic solution for fixation and transport of cytological samples.

1000 mL	CX-1L
---------	-------



CitoSpray

Solution for fixation of cytological smears in a spray bottle. Contains polyglycol which forms protective film when applied on cytological smear.

100 mL	CS-100
--------	--------



CitoSpray solution

Solution for fixation of cytological smears in a bottle, for refill of CitoSpray bottle.

1000 mL	CS-1L
---------	-------



Saccomanno's fixative

Solution for fixation and transport of cytological samples.

500 mL	SF-OT-500
1000 mL	SF-OT-1L
5000 mL	SF-OT-5L



Dehydrating and rehydrating agents

Alcoholic solutions and absolute alcohol are ideal for histocytology and cytology processing, storing and transporting samples. They are also used for dehydration and rehydration of tested samples.

Histanol 50

Denaturated 50% ethyl alcohol.

1000 mL	H50-1L
5000 mL	H50-5L
10000 mL	H50-10L



Histanol 70

Denaturated 70% ethyl alcohol.

1000 mL	H70-1L
5000 mL	H70-5L
10000 mL	H70-10L



Histanol 80

Denaturated 80% ethyl alcohol.

1000 mL	H80-1L
5000 mL	H80-5L
10000 mL	H80-10L



Histanol 95

Denaturated 95% ethyl alcohol.

1000 mL	H95-1L
5000 mL	H95-5L
10000 mL	H95-10L



Histanol 96

Denaturated 96% ethyl alcohol.

1000 mL	H96-1L
5000 mL	H96-5L
10000 mL	H96-10L



Histanol 100

Denaturated 100% ethyl alcohol.

1000 mL	H100-1L
5000 mL	H100-5L
10000 mL	H100-10L



Histanol EP 95

95% mixture of denaturated ethyl alcohol and isopropyl alcohol.

1000 mL	HEP95-1L
5000 mL	HEP95-5L
10000 mL	HEP95-10L



Histanol EP 100

100% mixture of denaturated ethyl alcohol and isopropyl alcohol.

1000 mL	HEP100-1L
5000 mL	HEP100-5L
10000 mL	HEP100-10L



Histanol EP 9010

Alcoholic solvent substituting standard ethanol series in processing, dewaxing and dehydrating.

1000 mL	HEP91-1L
5000 mL	HEP91-5L
10000 mL	HEP91-10L



Histanol IP

Isopropyl alcohol for use in microscopy.

1000 mL	HIP-1L
5000 mL	HIP-5L
10000 mL	HIP-10L



Histanol M

Methyl alcohol for use in microscopy.

1000 mL	HM-1L
5000 mL	HM-5L
10000 mL	HM-10L



Histanol G

Glycerol for use in microscopy.

1000 mL	HG-1L
5000 mL	HG-5L
10000 mL	HG-10L





CryoFix Gel

Embedding matrix for cryosectioning of tissue samples known as OCT medium (optimal cutting temperature). Efficient at temperatures of -10°C and below. Available in 5 different colors.

CryoFix Gel	100 mL	CF-100
CryoFix Gel pink	100 mL	CFR-100
CryoFix Gel blue	100 mL	CFP-100
CryoFix Gel green	100 mL	CFZ-100
CryoFix Gel yellow	100 mL	CFY-100



Cryo Spray

Medium in spray bottle for freezing histological samples.

200 mL 08-SPRAY-2



DeMold Spray

Solution in spray bottle for easy removal of paraffin tissue blocks from the mold.

100 mL DS-100



DeMold solution

Solution for refilling DeMold Spray bottle.

500 mL DS-500

1000 mL DS-1L



Tissue Soft

Solution for softening the tissue embedded in paraffin block before microtome sectioning.

500 mL TS-500

1000 mL TS-1L





BioGnost's synthetic BioMount media properties

PRODUCT NAME	SOLVENT	POLYMER	VISCOSITY @ 20°C (cSt, mPa)	REFRACTIVE INDEX @ 20°C (n _D)	USAGE IN AUTOMATED COVER SLIP MACHINES
BIOMOUNT DPX	xylene (BioClear)	S	850-950	1.5210-1.5230	-
BIOMOUNT DPX HIGH	xylene (BioClear)	S	1900-2100	1.5235-1.5255	-
BIOMOUNT DPX LOW	xylene (BioClear)	S	140-160	1.5115-1.5135	✓
BIOMOUNT DPX NEW*	xylene (BioClear)	S	200-300	1.5150-1.5180	✓
BIOMOUNT NEW	isoparaffin (BioClear New)	M	950-1050	1.4385-1.4395	-
BIOMOUNT NEW LOW	isoparaffin (BioClear New)	M	130-160	1.4345-1.4365	✓
BIOMOUNT	xylene (BioClear)	M	450-650	1.4920-1.4930	✓
BIOMOUNT HIGH	xylene (BioClear)	M	5000-5500	1.4900-1.4920	-
BIOMOUNT M	toluene	M	140-160	1.4900-1.4920	✓
BIOMOUNT C	xylene (BioClear)	M	5000-5500	1.4910-1.4925	-

Legend: M – methacrylate, S – styrene (polystyrene)
 * does not contain harmful dibutyl phthalate plasticizer
 ** contact us to find out how to decrease viscosity of a medium

Xylene and styrene-based media for permanent mounting

BioMount DPX

Moderate viscosity medium for permanent covering of microscope samples and mounting cover glass.

- 100 mL BM-100
- 250 mL BM-250
- 500 mL BM-500



BioMount DPX High

High viscosity medium for permanent covering of microscope samples and mounting cover glass.

- 100 mL BMH-100
- 250 mL BMH-250
- 500 mL BMH-500



BioMount DPX Low

Low viscosity medium for permanent covering of microscope samples and mounting cover glass. Can be used in automatic cover slip machines.

- 100 mL BML-100
- 250 mL BML-250
- 500 mL BML-500



BioMount DPX New

Low viscosity medium for permanent covering of microscope samples and mounting cover glass. Medium contains more environmentally friendly plasticizer instead of harmful dibutyl phthalate. Can be used in automatic cover slip machines.

- 100 mL BMLE-100
- 250 mL BMLE-250
- 500 mL BMLE-500



Methacrylate and xylene substitute-based media for permanent mounting

BioMount New

Moderate viscosity medium for permanent covering of microscope samples deparaffinized using xylene substitute-based medium, as well as xylene-based medium.

100 mL	BMN-100
250 mL	BMN-250
500 mL	BMN-500

BioMount New Low

Very low viscosity medium for permanent covering of microscope samples deparaffinized using xylene substitute, as well as xylene.

100 mL	BMNL-100
250 mL	BMNL-250
500 mL	BMNL-500



Methacrylate-based permanent mounting media

BioMount

Low viscosity xylene-based medium for permanent covering of microscope slides and mounting cover glass.

100 mL	BMT-100
250 mL	BMT-250
500 mL	BMT-500



BioMount M

Very low viscosity toluene-based medium for permanent covering of microscope samples and mounting cover glass.

100 mL	BMM-100
250 mL	BMM-250
500 mL	BMM-500



BioMount High

Very high viscosity xylene-based medium for permanent covering of microscope samples and mounting cover glass.

100 mL	BMTH-100
250 mL	BMTH-250
500 mL	BMTH-500





Additional permanent mounting media

BioMount C

Very high viscosity medium for permanent covering of microscope slides and mounting cover glass.

- 100 mL BMC-100
- 250 mL BMC-250
- 500 mL BMC-500

* excellent synthetic substitute for natural covering media, such as Canada Balsam



Water-based mounting media

PRODUCT NAME	SOLVENT	POLYMER	VISCOSITY @ 20°C (cSt, mPa)	REFRACTIVE INDEX @ 20°C (n _D)	USAGE IN AUTOMATED COVER SLIP MACHINES
BIOMOUNT AQUA	water	PVP	700-1000	1.4145-1.4205	-



BioMount Aqua

Water-based medium for covering microscope slides and mounting cover slips. With application in hematology, immunohistochemistry and immunocytochemistry with Fast-Red or AEC chromogens that require an aqueous mounting medium, but also with DAB chromogen.

- 30 mL BMA-30
- 100 mL BMA-100
- 250 mL BMA-250
- 500 mL BMA-500

Immersion oils

Seven different types of immersion oils for microscopical analysis

Immersion oils are essential for routine laboratory work, especially in hematology, cytology and microbiology. They are commonly used for high magnifications in order to secure high resolution and a clear image. There are several types of immersion oils that differ according to viscosity and certain properties required by different microscopy techniques. They all have a common refractive index of 1.5150, equal to the refractive index of glass.

BioGnost's Immersion oils properties

PRODUCT NAME	VISCOSITY @ 23°C (cSt, mPa)	COMBINING WITH OTHER TYPES OF OIL	TYPE OF MICROSCOPY
IMMERSION OIL	800-1300	Immersion oil type A Immersion oil type C	light
IMMERSION OIL TYPE A	100-300	Immersion oil Immersion oil type C	light
IMMERSION OIL TYPE C	1500-2200	Immersion oil Immersion oil type A	light
IMMERSION OIL TYPE FF	170	-	fluorescent
IMMERSION OIL TYPE 37*	1250**	-	light, for use at 37°C
IMMERSION OIL, TROPICAL GRADE***	1000-1500****	-	light, for use at 27-29°C

* refractive index n_D at 37°C = 1.5150 ** viscosity at 37°C *** refractive index n_D at tropical temperatures = 1.515 **** viscosity at 28°C

Immersion oil

Immersion oil for microscopy analysis with optimal viscosity of 800-1300 cSt.

10x10 mL	IU-1010
30 mL	IU-30
100 mL	IU-100
250 mL	IU-250
500 mL	IU-500

Immersion oil type A

Immersion oil for microscopy analysis with low viscosity of 100-300 cSt.

10x10 mL	IUA-1010
30 mL	IUA-30
100 mL	IUA-100
250 mL	IUA-250
500 mL	IUA-500





Immersion oil type C

Immersion oil for microscopy analysis with high viscosity of 1500-2200 cSt.

100 mL	IUC-100
250 mL	IUC-250
500 mL	IUC-500

* excellent synthetic substitute for natural immersion oils, such as Cedarwood oil



Immersion oil type FF

Immersion oil for sample analysis in fluorescent microscopy.

100 mL	IUF-100
250 mL	IUF-250
500 mL	IUF-500

Immersion oil type 37

Immersion oil for microscopy analysis with optimal refractive index and viscosity of 1250 cSt at 37°C.

100 mL	IU37-100
250 mL	IU37-250
500 mL	IU37-500



Immersion oil, tropical grade

Immersion oil for microscopy analysis in tropical climates with optimal refractive index at 27-29°C.

10x10 mL	IUT-1010
30 mL	IUT-30
100 mL	IUT-100
250 mL	IUT-250
500 mL	IUT-500



Cleaning solutions for microscope lens and slides

Clean Lens, solution 1

Solution according to Carl Zeiss formulation for cleaning microscope lens. Based on n-hexane and 2-propanol.

100 mL	CL1-OT-100
--------	------------



Clean Lens, solution 2

Solution for cleaning microscope lens and removing immersion oil from stained slide after using microscope. Based on diethyl ether and 2-propanol.

100 mL	CL2-OT-100
--------	------------



Clean Lens, solution 3

Solution for cleaning microscope lens and removing immersion oil from stained slide after using microscope. Based on petroleum ether and 2-propanol.

100 mL	CL3-OT-100
--------	------------



Standard histology staining reagents



Reagents for hematoxylin and eosin (HE) staining, the gold standard for demonstrating cellular and tissue structures, providing detailed anatomy and morphology of frozen and paraffin tissue sections, enabling clear visualization of nuclear, cytoplasmic and other cell structures.

Hematoxylin reagents

Nuclear stains for standard HE staining

Properties of BioGnost's Hematoxylin reagents

PRODUCT NAME	CYTOLOGY	HISTOLOGY	IHC	PROGRESSIVE METHOD	REGRESSIVE METHOD
HEMATOXYLIN H	-	✓	-	✓	✓
HEMATOXYLIN M	-	✓	✓	✓	-
HEMATOXYLIN ML	-	✓	✓	✓	✓
HEMATOXYLIN G1	✓	✓	✓	✓	-
HEMATOXYLIN G2	✓	✓	✓	✓	✓
HEMATOXYLIN G3	✓	✓	-	✓	✓
HEMATOXYLIN CAR	-	✓	✓	✓	-
HEMATOXYLIN COL	-	✓	✓	✓	✓
HEMATOXYLIN HP, PAP 1A	✓	-	-	✓	✓
HEMATOXYLIN HP, PAP 1B	✓	-	-	✓	✓
HEMATOXYLIN W KIT	-	✓	✓	✓	acid resistant
HEMATOXYLIN P.T.A.	-	✓	-	✓	-



Each of BioGnost's Hematoxylin is specially stabilized in order to maintain the reagent's reactivity for prolonged periods of time, as well as to maintain the quality of the reagent. Every staining provides a crystal clear and highly detailed image of the cell's nucleus, which in turn makes abnormal or pathological changes easier to detect.

Hematoxylin H

Modified hematoxylin according to Harris, most commonly used formulation in HE staining. Strong intensity reagent for progressive and regressive staining in histopathology.

100 mL	HEMH-OT-100
500 mL	HEMH-OT-500
1000 mL	HEMH-OT-1L
2500 mL	HEMH-OT-2.5L



Hematoxylin M

Modified hematoxylin according to Mayer. Weaker intensity reagent for progressive staining in histopathology and counterstaining in immunohistochemistry.

100 mL	HEMM-OT-100
500 mL	HEMM-OT-500
1000 mL	HEMM-OT-1L
2500 mL	HEMM-OT-2.5L



Hematoxylin ML

Modified hematoxylin according to Mayer-Lillie. Strong intensity new generation reagent for progressive and regressive staining in histopathology.

100 mL	HEMML-OT-100
500 mL	HEMML-OT-500
1000 mL	HEMML-OT-1L
2500 mL	HEMML-OT-2.5L



Hematoxylin G1

Hematoxylin according to Gill, ideal for staining goblet cells. New generation reagent for delicate progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG1-OT-100
500 mL	HEMG1-OT-500
1000 mL	HEMG1-OT-1L
2500 mL	HEMG1-OT-2.5L



Hematoxylin G2

Hematoxylin according to Gill, contains double concentration of hematoxylin compared to Hematoxylin G1. Moderate intensity new generation reagent for progressive and regressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG2-OT-100
500 mL	HEMG2-OT-500
1000 mL	HEMG2-OT-1L
2500 mL	HEMG2-OT-2.5L



Hematoxylin G3

Modified hematoxylin contains triple concentration of hematoxylin compared to Hematoxylin G1. Strong intensity new generation reagent for progressive and regressive staining in histopathology and cytology.

100 mL	HEMG3-OT-100
500 mL	HEMG3-OT-500
1000 mL	HEMG3-OT-1L
2500 mL	HEMG3-OT-2.5L



Hematoxylin W kit

Acid-resistant hematoxylin according to Weigert. Two-reagent kit that stains the nuclei blue-black, often a component of special staining kits for connective tissues.

2x100 mL	HEMW-K-100
2x500 mL	HEMW-K-500
2x1000 mL	HEMW-K-1L



Hematoxylin P.T.A.

Synonyms: Hematoxylin Phosphotungstic Acid, Mallory P.T.A.H. reagent. Simultaneously stains different types of tissue reddish-brown and blue. For differentiation between fibrin, smooth and cross-striated muscle fibers.

500 mL	HPTA-OT-500
1000 mL	HPTA-OT-1L

Hematoxylin CAR

Modified hematoxylin according to Carazzi for delicate nuclear staining without staining cytoplasmic components. Reagent for progressive staining of frozen sections in histopathology.

100 mL	HEMCA-OT-100
500 mL	HEMCA-OT-500
1000 mL	HEMCA-OT-1L

Hematoxylin COL

Modified hematoxylin according to Cole, contains iodine. Weaker intensity reagent for progressive and regressive staining in histopathology.

100 mL	HEMCO-OT-100
500 mL	HEMCO-OT-500
1000 mL	HEMCO-OT-1L





Hematoxylin Instant kit

Two-component kit that contains ready-to-use powders for preparation of hematoxylin reagent for progressive and regressive nuclear staining in histopathology and cytology.

for 1000 mL HEMI-K-1L
for 6000 mL HEMI-K-6L

Kit for rapid HE staining of frozen and paraffin-embedded sections

Ready-to-use eight-reagent kit (in 20 containers that can be used as staining jars) for rapid HE staining of frozen and paraffin tissue sections in histopathology. Contains xylene substitute as clearing agent and xylene substitute-based medium for permanent coverslipping.

for 100 tests HE-RTU-100T



ErioGnost reagent

Acid fast reagent for nuclear staining, comparable to hematoxylin staining. Eco-friendly synthetic replacement for hematoxylin reagents.

100 mL ERS-OT-100
500 mL ERS-OT-500
1000 mL ERS-OT-1L



Differentiation and bluing reagents for histology and cytology

Differentiation of the nucleus enables clear visualization of nuclear structures in regressive procedure, while the bluing reagent turns the nucleus blue.

Acid alcohol

Reagent for enabling crisp visualization of nuclear chromatin detail and removing of excess background dye in histological samples after regressive staining with hematoxylin.

1000 mL KA-OT-1L
2500 mL KA-OT-2.5L



Scott's solution

Buffered reagent with potassium carbonate and magnesium sulphate for bluing of nuclear chromatin and nuclear membranes.

1000 mL SC-OT-1L
2500 mL SC-OT-2.5L

HCL Pap reagent

Aqueous reagent for enabling crisp visualization of nuclear chromatin detail and removing of excess background dye in cytological after regressive staining with hematoxylin.

1000 mL HCLPAP-OT-1L
2500 mL HCLPAP-OT-2.5L

Bluing reagent

Blue colored buffered reagent with lithium carbonate for bluing of nuclear chromatin and nuclear membranes.

1000 mL BR-OT-1L
2500 mL BR-OT-2.5L

NEW



BioBluing buffer

Buffered reagent for nuclear bluing in standard HE staining. Ensures optimal staining and proper cellular color staining. Because of alkaline pH, it turns the hematoxylin blue. Premixed, ready-to-use and tinted blue for easy identification. Utilizes a buffer system to prevent pH fluctuation.

1000 mL BB-OT-1L
2500 mL BB-OT-2.5L



Hem Diff

Differentiation reagent in HE modified regressive staining method. Produces crisp nuclear chromatin detail and eliminates undesirable background staining. Premixed and ready-to-use. Non-hazardous. Used for differentiation of Hematoxylin G2, G3 and ML

1000 mL HD-OT-1L
2500 mL HD-OT-2.5L



Hem Diff Strong

Differentiation reagent in HE regressive staining method. Provides excellent differentiation between nuclear and non-nuclear structures. Premixed and ready-to-use. Based on acid alcohol. Used for differentiation of Hematoxylin H.

1000 mL HDS-OT-1L
2500 mL HDS-OT-2.5L

Eosin reagents

Cytoplasmic stains for standard HE staining

Eosin Y (yellowish eosin) solutions are commonly used as a counterstain to hematoxylin in the standard HE staining method. Eosin Y is an anionic dye that stains basic cell components (such as cytoplasm, collagen, muscle fibers and erythrocytes) bright red.



Eosin Y aqueous solutions

Eosin Y aqueous solutions for cytoplasmic counterstaining.

Eosin Y 0.2% aqueous	1000 mL	EOY-02-OT-1L	
	2500 mL	EOY-02-OT-2.5L	
Eosin Y 0.5% aqueous	1000 mL	EOY-05-OT-1L	
	2500 mL	EOY-05-OT-2.5L	
Eosin Y 1% aqueous	1000 mL	EOY-10-OT-1L	
	2500 mL	EOY-10-OT-2.5L	
Eosin Y 2% aqueous	1000 mL	EOY-20-OT-1L	
	2500 mL	EOY-20-OT-2.5L	

Eosin Y alcoholic solutions

Eosin Y alcoholic solutions for cytoplasmic counterstaining.

Eosin Y 0.5% alcoholic	1000 mL	EOYA-05-OT-1L	
	2500 mL	EOYA-05-OT-2.5L	
Eosin Y 1% alcoholic	1000 mL	EOYA-10-OT-1L	
	2500 mL	EOYA-10-OT-2.5L	
Eosin Y 2% alcoholic	1000 mL	EOYA-20-OT-1L	
	2500 mL	EOYA-20-OT-2.5L	

Eosin Contrast

Modified alcoholic solution for cytoplasmic counterstaining. The reagent contains Eosin Y and Phloxine B for enhanced counterstaining effect.

1000 mL	EOYK-OT-1L	
2500 mL	EOYK-OT-2.5L	

Eosin Contrast PLUS

Modified alcoholic solution for intensive cytoplasmic counterstaining. Reagent contains Eosin Y, Phloxine B, and Biebrich Scarlet dyes for additional counterstaining effect.

1000 mL	EOYKP-OT-1L	
2500 mL	EOYKP-OT-2.5L	

Eosin Y instant

Mixture of powders for preparation of eosin reagent in aqueous or alcoholic medium.

Eosin Y Instant, aqueous	for 1000 mL	EOYI-AQ-1L
Eosin Y Instant, alcoholic	for 1000 mL	EOYI-ALC-1L





Cytology and hematology staining reagents

Cytology and hematology staining kits contain reagents for fixation, staining and buffering cytology and hematology samples, as well as cytology smears of hematopoietic organs, providing precise distinction between cellular structures. Cytochemical staining kits contain reagents for cytochemical diagnosis of leukemia using bone marrow or whole blood smears, and also for staining and determining specific cell components in blood samples.

Papanicolaou (Pap) reagents

Nuclear and cytoplasmic reagents for Pap staining

Due to their simplicity of use and exceptional precision in diagnostics, staining reagents according to Papanicolaou have substantially contributed to decreasing occurrence of cervical cancer. By using a simple staining protocol, BioGnost's Pap reagents enable excellent differentiation between cellular structures.



Hematoxylin HP, Pap 1A

Modified Harris hematoxylin solution for nuclear staining according to Papanicolaou. Strong intensity reagent for progressive and regressive staining in exfoliative cytology.

100 mL	HEMHP-OT-100
500 mL	HEMHP-OT-500
1000 mL	HEMHP-OT-1L
2500 mL	HEMHP-OT-2.5L

Hematoxylin HP, Pap 1B

Hematoxylin modified according to Harris for intensive nuclear staining according to Papanicolaou. Reagent for progressive and regressive staining in exfoliative cytology, with formulation stronger than Hematoxylin HP, Pap 1A.

100 mL	HEMHP1B-OT-100
500 mL	HEMHP1B-OT-500
1000 mL	HEMHP1B-OT-1L
2500 mL	HEMHP1B-OT-2.5L

OG-6 reagent, Pap 2A

Monochromatic reagent for counterstaining mature and keratinized squamous cell. Contains Orange G dye that stains the cytoplasm orange.

100 mL	OG6-OT-100
500 mL	OG6-OT-500
1000 mL	OG6-OT-1L
2500 mL	OG6-OT-2.5L

Orange II reagent, Pap 2B

Monochromatic reagent for counterstaining mature and keratinized squamous cells. Contains Orange II dye that stains the cytoplasm intense red.

100 mL	OR2-OT-100
500 mL	OR2-OT-500
1000 mL	OR2-OT-1L
2500 mL	OR2-OT-2.5L

EA 31 reagent, Pap 3A

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **gynecology** samples in cytology, contains the largest amount of green dye compared to other EA reagents. Stains the cytoplasm blue-green and pink, with emphasis on green and pink.

100 mL	EA31-OT-100
500 mL	EA31-OT-500
1000 mL	EA31-OT-1L
2500 mL	EA31-OT-2.5L

EA 50 reagent, Pap 3B

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **gynecology** samples in cytology, stains the cytoplasm blue-green and pink, with emphasis on blue and pink.

100 mL	EA50-OT-100
500 mL	EA50-OT-500
1000 mL	EA50-OT-1L
2500 mL	EA50-OT-2.5L

EA 65 reagent, Pap 3C

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **non-gynecology** samples in cytology, stains the cytoplasm red.

100 mL	EA65C-OT-100
500 mL	EA65C-OT-500
1000 mL	EA65C-OT-1L
2500 mL	EA65C-OT-2.5L

EA 65 reagent, Pap 3D

Polychromatic reagent for staining and differentiation of squamous cells. Counterstain for **non-gynecology** samples in cytology, stains the cytoplasm light blue-green and pink.

100 mL	EA65D-OT-100
500 mL	EA65D-OT-500
1000 mL	EA65D-OT-1L
2500 mL	EA65D-OT-2.5L

Rapid Pap staining kit

Ready-to-use eight-reagent kit (in 20 containers that can be used as staining jars) for rapid progressive gynecology and non-gynecology cytological samples. Contains xylene substitute as clearing agent and xylene substitute-based medium for permanent covering of samples.

for 100 tests	PAP-100T
---------------	----------

OG-EA31 Pap reagent

One-step polychromatic reagent used as counterstain for gynecology samples in cytology. Enables rapid staining and differentiation between epithelial cells whose cytoplasm is stained green and pink. It is not necessary to use OG-6, Pap 2A reagent during staining because it stains cytoplasm of keratinized cell orange at the same time.

100 mL	OGEA31-OT-100
500 mL	OGEA31-OT-500
1000 mL	OGEA31-OT-1L
2500 mL	OGEA31-OT-2.5L

OG-EA50 Pap reagent

One-step polychromatic reagent used as counterstain for gynecology samples in cytology. Enables rapid staining and differentiation between epithelial cells whose cytoplasm is stained blue and pink. It is not necessary to use OG-6, Pap 2A reagent during staining because it stains cytoplasm of keratinized cell orange at the same time.

100 mL	OGEA50-OT-100
500 mL	OGEA50-OT-500
1000 mL	OGEA50-OT-1L
2500 mL	OGEA50-OT-2.5L

Shorr's solution

Enables clear differentiation of eosinophilic and cyanophilic cells of cervical epithelium. It is also used for a better visualization of sperm morphology.

100 mL	SH-OT-100
500 mL	SH-OT-500
1000 mL	SH-OT-1L
2500 mL	SH-OT-2.5L

Hematoxylin G1

Hematoxylin according to Gill, ideal for staining goblet cells. New generation reagent for delicate progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG1-OT-100
500 mL	HEMG1-OT-500
1000 mL	HEMG1-OT-1L
2500 mL	HEMG1-OT-2.5L

Hematoxylin G2


Hematoxylin according to Gill, contains double concentration of hematoxylin compared to Hematoxylin G1. Moderate intensity new generation reagent for progressive staining in histopathology, cytology and for counterstaining in immunohistochemistry.

100 mL	HEMG2-OT-100
500 mL	HEMG2-OT-500
1000 mL	HEMG2-OT-1L
2500 mL	HEMG2-OT-2.5L

Hematoxylin G3

Modified hematoxylin contains triple concentration of hematoxylin compared to Hematoxylin G1. Strong intensity new generation reagent for progressive staining in histopathology and cytology.

100 mL	HEMG3-OT-100
500 mL	HEMG3-OT-500
1000 mL	HEMG3-OT-1L
2500 mL	HEMG3-OT-2.5L





Rapid and standard hematology reagents and kits

Bio-Diff kits

Rapid staining solutions for hematology, cytology and histology

Rapid diagnostic kits that provide precise staining results in a short period of time, such as results provided by May-Gruenwald Giemsa (MGG) staining method. Each component of Bio-Diff and Bio-Diff RTU kits is stabilized and prepared according to the highest standards.

Bio-Diff kit

Three-reagent kit that contains fixative agent, red and blue components for fast and effective staining. Each kit contains buffer tablets for consistent staining results.

3x100 mL	BD-K-100
3x500 mL	BD-K-500
3x1000 mL	BD-K-1L
3x2500 mL	BD-K-2.5L



Bio-Diff 1 reagent

Fixative agent, component of Bio-Diff kit.

100 mL	BD1-OT-100
500 mL	BD1-OT-500
1000 mL	BD1-OT-1L
2500 mL	BD1-OT-2.5L



Bio-Diff 2 reagent

Buffered red component of Bio-Diff kit.

100 mL	BD2-OT-100
500 mL	BD2-OT-500
1000 mL	BD2-OT-1L
2500 mL	BD2-OT-2.5L

Bio-Diff 3 reagent

Buffered blue component of Bio-Diff kit.

100 mL	BD3-OT-100
500 mL	BD3-OT-500
1000 mL	BD3-OT-1L
2500 mL	BD3-OT-2.5L

STAINING DONE IN 14 SECONDS*



* Depending on the type of sample.

Bio-Diff RTU kit

Ready-to-Use three-reagent kit with reagents in containers that can be used as staining jars. Kit contains fixative agent, red and blue components for fast and effective staining and buffer tablet for consistent staining results.

3x100 mL BD-RTU-100



Bio-Diff 1 RTU reagent

Fixative component Bio-Diff RTU kit.

100 mL BD1-RTU



Bio-Diff 2 RTU reagent

Buffered red component of Bio-Diff RTU kit.

100 mL BD2-RTU

Bio-Diff 3 RTU reagent

Buffered blue component of Bio-Diff RTU kit.

100 mL BD3-RTU

STAINING DONE IN
14 SECONDS*



* depending on the type of sample

Buffer solution, pH 7.2

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

1000 mL PUF72-OT-1L
2500 mL PUF72-OT-2.5L
5000 mL PUF72-OT-5L

Buffer tablets pH 7.2

Phosphate buffer tablets for preparing buffer solutions used in hematology and histology. One buffer tablet is sufficient for preparation of 1000 mL of phosphate buffer solution.

50 pcs PT-72-50
100 pcs PT-72-100



Use buffer solution pH 6.8 for hematological and cytological diagnostics. Use buffer solution pH 7.2 for the detection of blood parasites.



May-Gruenwald and Giemsa reagents

Romanowsky staining solutions for hematology, cytology and histology

One of the most commonly used Pappenheim methods is May-Gruenwald Giemsa (MGG) for staining hematology and cytology materials, as well as cytology smears of hematopoietic organs. Giemsa solution is also used for staining histology sections.

May-Gruenwald solution

Polychromatic fixative solution containing Eosin Y, Methylene Blue and azure dyes. For staining in hematology, cytology and staining sections of hematopoietic organs in histopathology.

100 mL	MG-OT-100
10x100 mL	MG-OT-110
500 mL	MG-OT-500
1000 mL	MG-OT-1L
2500 mL	MG-OT-2.5L

Giemsa solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For use in hematology, cytology, histology and for staining section samples of hematopoietic organs.

100 mL	GM-OT-100
10x100 mL	GM-OT-110
500 mL	GM-OT-500
1000 mL	GM-OT-1L
2500 mL	GM-OT-2.5L

Buffer solution, pH 6.4

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

1000 mL	PUF64-OT-1L
2500 mL	PUF64-OT-2.5L
5000 mL	PUF64-OT-5L

Buffer solution, pH 6.8

Phosphate buffer solution, molarity 6 mM for use in hematology and histology.

1000 mL	PUF68-OT-1L
2500 mL	PUF68-OT-2.5L
5000 mL	PUF68-OT-5L

Buffer tablets pH 6.8

Phosphate buffer tablets for preparing solutions used in hematology and histology. One buffer tablet is sufficient for preparation of 1000 mL of phosphate buffer solution.

50 pcs	PT-68-50
100 pcs	PT-68-100

Use buffer solution pH 6.8 for hematological and cytological diagnostics. Use buffer solution pH 7.2 for the detection of blood parasites.



Additional staining solutions for hematology and cytology



Brilliant Cresyl Blue, solution

Solution for staining reticulocytes in hematology.

100 mL	BCB-OT-100
500 mL	BCB-OT-500
1000 mL	BCB-OT-1L

Leishman's solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For staining in hematology and clinical cytology.

100 mL	LE-OT-100
500 mL	LE-OT-500
1000 mL	LE-OT-1L

Wright's solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. For staining in hematology, cytology and cytogenetics.

100 mL	WR-OT-100
500 mL	WR-OT-500
1000 mL	WR-OT-1L

Field kit

Ready to use two-reagent kit for rapid and efficient staining and detection of parasites in hematology samples. Kit for 100 tests contains reagents in containers that can be used as staining jars.

for 100 tests	FD-100T
2x500 mL	FD-K-500

Wright-Giemsa solution

Polychromatic solution of Eosin Y, Methylene Blue and azure dyes. Modified Romanowsky stain for hematology staining.

100 mL	WRGM-OT-100
500 mL	WRGM-OT-500
1000 mL	WRGM-OT-1L
2500 mL	WRGM-OT-2.5L

Hayem's solution

Solution for manual counting of erythrocytes.

100 mL	HY-OT-100
500 mL	HY-OT-500
1000 mL	HY-OT-1L

Tuerk's solution

Solution for manual counting of leukocytes.

100 mL	TU-OT-100
500 mL	TU-OT-500
1000 mL	TU-OT-1L

Bio-Hansel solution NEW

Polychromatic solution resistant to an acidic environment used for fast and accurate identification of eosinophilic granules in hematological and cytological samples.

100 mL	BHO-OT-100
--------	------------

CytoGnost kit

Three-reagent kit for manual counting of human blood components.

3 x 100 mL	HCG-K-100
3 x 500 mL	HCG-K-500
3 x 1000 mL	HCG-K-1L

ThromboGnost solution

Solution for manual counting of thrombocytes.

100 mL	THR-OT-100
500 mL	THR-OT-500
1000 mL	THR-OT-1L



LEUKOGNOST

Cytochemical staining kits and ancillary reagents

Cytochemical staining kits

Enzymocytochemical blood smear and bone marrow staining of sections used to determine location and activity of cellular structures and enzyme systems in leukemia cell cytoplasm. Known as a routine method for diagnosis of hematological diseases and leukemia classification.

LeukoGnost SPE

Kit for detection of specific esterase activity (Leder stain, chloroacetate esterase) in leukocytes. Can be applied on formalin-fixed, paraffin-embedded tissue.

for at least 100 tests LKG-SPE



LeukoGnost ALP

Kit for detection of alkaline phosphatase activity in leukocytes.

for at least 100 tests LKG-ALP



LeukoGnost MPO

Kit for detection of myeloperoxidase activity in leukocytes.

for at least 100 tests LKG-MPO



LeukoGnost PAS

Kit for detection of the periodic acid-Schiff reaction in leukocytes.

for at least 100 tests LKG-PAS



LeukoGnost NSE

Kit for detection of non-specific esterase activity in leukocytes.

for at least 100 tests LKG-NSE



LeukoGnost SPENSE

Kit for simultaneous detection of specific and non-specific esterase activity in leukocytes.

for at least 50 tests LKG-SPENSE



LeukoGnost ACP

Kit for detection of acid phosphatase activity in leukocytes.

for at least 50 tests LKG-ACP



LeukoGnost PERLS

LeukoGnost PERLS (Prussian blue) kit for detection of free ferric (Fe³⁺) ions in blood and bone marrow. For application on tissue cells use HemoGnost PERLS kit.

for 100 tests LKG-PERLS

Sudan Black B kit

Four-reagent kit for staining neutrophil granules in hematological smears.

for 350-400 tests SBBH-K-100



Sudan Black B ECO

Phenol-free three-reagent kit for staining neutrophil granules in hematological smears.

for 100 tests SBBE-100T





Bacteriology and mycology staining reagents

Reagents for rapid and easy staining in microbiology allowing optimal differential analysis of bacteria. Phenol-free staining kits offer environment-friendly alternative without compromising quality of staining. Reagents for fungal analysis enable visualization of fungal structures (otherwise difficult to observe) and samples preservation.

BioGram kits and reagents

Kits for identification of bacteria according to Gram

Differential staining of bacteria according to Gram is widely known and used method in microbiology. BioGram 4 kit uses standard method to provide optimal staining and identification of Gram-positive and Gram-negative bacteria. BioGram ECO kit is a phenol-free, eco-friendly alternative to BioGram 4 kit with equal staining results.

BioGram 4 kit

Four-reagent kit for identification of bacteria according to Gram. Kit contains Gram Crystal Violet 1% solution, stabilized Gram Lugol solution, double amount of Gram Decolorizer solution 2 and Gram Safranin solution as counterstain.

for 100 tests	BGR4-100T
5x100 mL	BGR4-K-100
5x250 mL	BGR4-K-250
5x500 mL	BGR4-K-500

Gram Crystal Violet 1% solution

Primary stain reagent for Gram-positive bacteria. Crystal Violet is also known as Gentian Violet and Methyl Violet 10B.

100 mL	GC1-OT-100
250 mL	GC1-OT-250
500 mL	GC1-OT-500
1000 mL	GC1-OT-1L
2500 mL	GC1-OT-2.5L

Gram Lugol solution, stabilized

Solution contains polyvinylpyrrolidone iodine (PVP-I).

100 mL	GLS-OT-100
250 mL	GLS-OT-250
500 mL	GLS-OT-500
1000 mL	GLS-OT-1L
2500 mL	GLS-OT-2.5L

Gram Decolorizer solution 2

Medium speed decolorizer used after stabilized iodine solution.

100 mL	GD2-OT-100
250 mL	GD2-OT-250
500 mL	GD2-OT-500
1000 mL	GD2-OT-1L
2500 mL	GD2-OT-2.5L

Gram Safranin solution

Counterstain for Gram-negative bacteria, contains Safranin O dye.

100 mL	GSF-OT-100
250 mL	GSF-OT-250
500 mL	GSF-OT-500
1000 mL	GSF-OT-1L
2500 mL	GSF-OT-2.5L

BioGram ECO kit PHENOL-FREE

Four-reagent phenol-free kit for identification of bacteria according to Gram. Kit contains Gram Crystal violet, phenol-free reagent, Gram Sodium hydrogencarbonate, solution, stabilized Gram Lugol solution, double amount of Gram Decolorizer solution 2 and Gram Safranin solution as counterstain.

2x50 mL+4x100 mL	BGRE-K-100
2x250 mL+4x500 mL	BGRE-K-500

Gram Crystal Violet, phenol-free reagent

Primary stain reagent for Gram-positive bacteria, doesn't contain phenol. Crystal Violet is also known as Gentian Violet and Methyl Violet 10B.

100 mL	GCF-OT-100
250 mL	GCF-OT-250
500 mL	GCF-OT-500
1000 mL	GCF-OT-1L
2500 mL	GCF-OT-2.5L

Gram Sodium hydrogencarbonate, solution

Solution of sodium hydrogencarbonate for use with BioGram ECO kit.

100 mL	GNHK-OT-100
250 mL	GNHK-OT-250
500 mL	GNHK-OT-500
1000 mL	GNHK-OT-1L
2500 mL	GNHK-OT-2.5L

Additional bacteriology staining solutions according to Gram

Gram Lugol solution

Non-stabilized Lugol solution.

100 mL	GL-OT-100
250 mL	GL-OT-250
500 mL	GL-OT-500
1000 mL	GL-OT-1L

Gram Decolorizer solution 1

Slow acting decolorizer solution used after non-stabilized iodine solution.

100 mL	GD1-OT-100
250 mL	GD1-OT-250
500 mL	GD1-OT-500
1000 mL	GD1-OT-1L

Gram Decolorizer solution 3

Fast acting solution used with automated stainers.

100 mL	GD3-OT-100
250 mL	GD3-OT-250
500 mL	GD3-OT-500
1000 mL	GD3-OT-1L

Gram Safranin 5x concentrated

Concentrated counterstain solution of Safranin O dye.

For preparation of 500 mL of Safranin O counterstain solution	100 mL	SO05X-OT-100
For preparation of 2500 mL of Safranin O counterstain solution	500 mL	SO05X-OT-500
For preparation of 5000 mL of Safranin O counterstain solution	1000 mL	SO05X-OT-1L

Carbol Fuchsin Gram 10x concentrated

Concentrated Carbol Fuchsin solution for preparation of counterstain solution for modified staining. Also known as concentrated Gram Fuchsin solution.

For preparation of 1000 mL of Carbol Fuchsin Gram counterstain solution	100 mL	CFGR10-OT-100
For preparation of 5000 mL of Carbol Fuchsin Gram counterstain solution	500 mL	CFGR10-OT-500
For preparation of 10000 mL of Carbol Fuchsin Gram counterstain solution	1000 mL	CFGR10-OT-1L

Additional iodine solutions

Lugol's solution

Classic formulation of aqueous solution of iodine and potassium iodide.

100 mL	LUG-OT-100
250 mL	LUG-OT-250
500 mL	LUG-OT-500

Lugol's C solution

Aqueous solution of iodine and potassium iodide, enhanced formulation with stronger staining (compared to classic Lugol solution).

100 mL	LUGC-OT-100
250 mL	LUGC-OT-250
500 mL	LUGC-OT-500

Lugol's solution, acid

Aqueous iodine and potassium iodide solution with added acid.

100 mL	LUGK-OT-100
250 mL	LUGK-OT-250
500 mL	LUGK-OT-500



TB-Stain kits and reagents

Kits for staining acid-fast bacteria according to Ziehl-Neelsen

Most commonly known method for detecting TB bacteria according to Ziehl-Neelsen. This method enables rapid and easy staining of acid-fast bacterial strains, such as *Mycobacteria*, *Nocardia*, *Rhodococcus*, *Dietzia* and *Gordonia*. TB-Stain ECO kit is a phenol-free, environment-friendly alternative to TB-Stain Hot kit.

TB-Stain Hot kit

Three-reagent kit for staining acid-fast bacteria. Contains TB Carbol Fuchsin reagent, double amount of TB Decolorizer and Methylene Blue Loeffler's reagent as counterstain.

for 100 tests	TBH-100T
4x100 mL	TBH-K-100
4x250 mL	TBH-K-250
4x500 mL	TBH-K-500



TB Carbol Fuchsin reagent

Synonym: Ziehl-Neelsen reagent. Primary stain reagent for acid-fast bacteria used in TB-Stain Hot, TB-Stain Cold and TB-Stain Quick kits.

100 mL	TBC-OT-100
250 mL	TBC-OT-250
500 mL	TBC-OT-500
1000 mL	TBC-OT-1L
2500 mL	TBC-OT-2.5L



TB-Stain ECO kit PHENOL-FREE

Three-reagent phenol-free kit for staining acid-fast bacteria. Contains TB-Fuchsin reagent, double amount of TB Decolorizer and Methylene Blue Loeffler's reagent as counterstain.

4x100 mL	TBE-K-100
4x250 mL	TBE-K-250
4x500 mL	TBE-K-500



TB Decolorizer

For use in TB-Stain Hot and TB-Stain Cold kits.

100 mL	TBD-OT-100
250 mL	TBD-OT-250
500 mL	TBD-OT-500
1000 mL	TBD-OT-1L
2500 mL	TBD-OT-2.5L



Methylene Blue Loeffler's reagent

Blue counterstain for TB-Stain Hot kit.

100 mL	MBL-OT-100
250 mL	MBL-OT-250
500 mL	MBL-OT-500
1000 mL	MBL-OT-1L
2500 mL	MBL-OT-2.5L

TB Fuchsin reagent

Primary stain phenol-free reagent for acid-fast bacteria used in TB-Stain ECO kit.

100 mL	TBFR-OT-100
250 mL	TBFR-OT-250
500 mL	TBFR-OT-500
1000 mL	TBFR-OT-1L
2500 mL	TBFR-OT-2.5L

Additional TB-Stain Hot kit staining solution

Methylene Blue 10x concentrated

Concentrated counterstain solution of Methylene Blue dye for use in microbiology.

For preparation of 1000 mL of Methylene Blue counterstain solution	100 mL	MB10X-OT-100
--	--------	--------------

For preparation of 5000 mL of Methylene Blue counterstain solution	500 mL	MB10X-OT-500
--	--------	--------------

For preparation of 10000 mL of Methylene Blue counterstain solution	1000 mL	MB10X-OT-1L
---	---------	-------------





Kits for detection of acid-fast bacteria using fluorescence microscopy according to modified Truant method

TB-Stain Fluorescent kit

Three-reagent kit for detection of acid-fast bacteria using fluorescence microscopy. Contains TB Auramine-Rhodamine reagent, double amount of TB Decolorizer Fluorescent and TB Permanganate reagent as counterstain.

for 100 tests	TBF-100T
4x100 mL	TBF-K-100
4x250 mL	TBF-K-250
4x500 mL	TBF-K-500



TB-Stain Auramine O kit

Three-reagent kit for staining acid-fast bacteria using fluorescence method. Contains TB Auramine O reagent, double amount of TB Decolorizer Fluorescent and counterstain of TB Permanganate reagent.

4x100 mL	TBAO-K-100
4x250 mL	TBAO-K-250
4x500 mL	TBAO-K-500



TB Auramine O reagent

Primary stain for use in TB Stain Auramine O kit.

100 mL	TBAO-OT-100
250 mL	TBAO-OT-250
500 mL	TBAO-OT-500
1000 mL	TBAO-OT-1L
2500 mL	TBAO-OT-2.5L



TB Auramine-Rhodamine reagent

Primary stain reagent for use in TB-Stain Fluorescent kit.

100 mL	TBA-OT-100
250 mL	TBA-OT-250
500 mL	TBA-OT-500
1000 mL	TBA-OT-1L
2500 mL	TBA-OT-2.5L



TB Decolorizer Fluorescent

For use in TB-Stain Fluorescent and TB-Stain Auramine O kits.

100 mL	TBF-OT-100
250 mL	TBF-OT-250
500 mL	TBF-OT-500
1000 mL	TBF-OT-1L
2500 mL	TBF-OT-2.5L



TB Permanganate reagent

Counterstain for use in TB-Stain Fluorescent kit and TB-Stain Auramine O kits.

100 mL	TBP-OT-100
250 mL	TBP-OT-250
500 mL	TBP-OT-500
1000 mL	TBP-OT-1L
2500 mL	TBP-OT-2.5L





Special staining kits for histology and cytology

Special kits for histology and cytology enable clear visualization of specific structures and cellular components. All of the reagents are easy to use and precisely formulated to maximize usability and to provide optimal results for the user.

Classification of special staining kits according to use:

MUSCLE AND/OR CONNECTIVE TISSUE

- A.F.O.G kit
- Azan Trichrome kit
- Elastica-Van Gieson kit
- Gomori Trichrome kit
- H.B.F.P. kit
- Hematoxylin P.T.A. kit
- Hematoxylin W kit
- Ladewig's Trichrome kit
- Mallory Trichrome kit
- Martius Scarlet Blue (MSB) kit
- Masson Trichrome kit
- Masson-Goldner Trichrome kit
- Movat kit
- Orcein kit
- Paraldehyde Fuchsin kit
- P.A.S.M. / Jones kit
- Picro-Sirius Red kit
- Safranin O kit
- Reticulin kit
- Reticulin Contrast kit
- Van Gieson Trichrome kit
- Verhoeff kit
- Weigert-Van Gieson kit

ENZYME ACTIVITY

- LeukoGnost SPE

FUNGI, BACTERIA AND PARASITES

- Alcian Yellow Toluidine Blue kit
- BioGram Histo kit
- Field kit
- Giemsa HP kit
- Grocott kit
- M.I.F. kit
- TB-Stain Histo kit
- Warthin Starry kit

LIPIDS

- Oil Red O kit
- Sudan Black B Lipid kit

NUCLEI AND NUCLEIC ACIDS

- Feulgen kit

PIGMENTS AND MINERAL DEPOSITS

- Fouchet-Van Gieson kit
- Grimelius kit
- HemoGnost Perls kit
- Masson Fontana kit
- Orcein kit
- Perls-Van Gieson kit
- Rhodanine kit
- Von Kossa kit

CYTOLOGY

- Bio-Diff RTU kit
- Eosin and nigrosin vital kit
- Sperm-Diff RTU kit
- UriGnost S kit
- UriGnost SM kit

CARBOHYDRATES


- Alcian Blue - P.A.S. kit
- Alcian Blue pH 1.0 kit
- Alcian Blue pH 2.5 kit
- Colloidal Iron kit
- Congo Red Highman kit
- Congo Red Puchtler kit
- Mucicarmine kit
- P.A.S. kit
- P.A.S. Diastase kit

CRYOSTAT

- Gomori Trichrome kit
- Oil Red O kit
- Orcein kit
- Sudan Black B Lipid kit

CENTRAL NERVOUS SYSTEM


- Bielschowsky kit
- Luxol Fast Blue kit



A.F.O.G. kit

Six-reagent Acid Euchsin Orange G kit for selective staining of glomerular protein deposits and collagen in kidney biopsies. Nuclear stain is obtained with Weigert ferric hematoxylin, cytoplasm with Orange G and highly selective collagen stain with Aniline blue.


for 100 tests AFOG-100T
6x100 mL AFOG-K-100



Alcian Blue - P.A.S. kit

Seven-reagent Alcian Blue - Periodic Acid-Schiff kit for staining acid mucopolysaccharides according to Mowry. Enables differentiation between acid mucins (stained light blue) and neutral mucins, glycogens and glycoproteins (stained magenta).


for 100 tests AB-100T
7x100 mL AB-K-100
7x500 mL AB-K-500



Alcian Blue pH 1.0 kit

Three-reagent kit for specific staining of heavily sulfated mucopolysaccharides blue. Slides are counterstained with Nuclear Fast Red (Kernechtrot) reagent to fully detect the presence of Alcian Blue positive staining.


for 100 tests AB10-100T
3x100 mL AB10-K-100



Alcian Blue pH 2.5 kit

Three-reagent kit for staining sulfated and carboxylated acid mucopolysaccharides and sulfated and carboxylated sialomucins blue. Slides are counterstained with Nuclear Fast Red (Kernechtrot) reagent to fully detect the presence of Alcian Blue positive staining.


for 100 tests AB25-100T
3x100 mL AB25-K-100



Alcian Yellow Toluidine Blue kit

Six-reagent kit for staining *Helicobacter pylori* in gastric tissue sections. This method is one of the most popular non-silver method for staining of *H. pylori*, where bacteria are stained blue in contrast to yellow mucins.

for 100 tests AYTB-100T




Azan Trichrome kit

Five-reagent kit for connective tissue staining according to Mallory. Used for visualization of muscle fibers, collagen, glial cells, kidney glomerular stroma, erythrocytes, reticulum and nuclear chromatin in histologic sections.

for 100 tests AZT-100T
5x250 mL AZT-K-250






Elastica-Van Gieson kit

Four-reagent kit for staining elastic fibers and differentiation between elastic tissue, collagen and other types of connective tissue. The rapid method enables a satisfactory result with shorter section staining time.


for 100 tests WVGB-100T
4x100 mL WVGB-K-100



Eosin and nigrosine vital kit

Two-reagent kit containing separate dyes for rapid detection of sperm vitality and simple visualization of dead and living sperm cells. The nigrosin stain provides dark background for easier recognition of both viable and non-viable spermatozoa.


2x30 mL ENVK-K-30
2x100 mL ENVK-K-100



Feulgen kit

Five-reagent DNA staining kit according to Feulgen. For use in semiquantitative DNA determination in histological and cytological samples. Specimen is first treated with hydrochloric acid creating an aldehyde group of DNA that can be visualized by Schiff (BioSchiff) reagent. This reaction is specific for nuclear DNA.


for 100 tests FE-100T
6x100 mL FE-K-100



Field kit

Ready-to-use two-reagent kit for rapid and efficient staining and detection of parasites in hematology samples. Primarily used for staining thin and thick blood smears (dense drop) for purpose of diagnosing blood parasites. Reagents are stored in containers that can be used as staining jars.


for 100 tests FD-100T
2x500 mL FD-K-500



Fouchet-Van Gieson kit

Three-reagent kit for visualization of bilirubin and collagen according to Kutllick. Bilirubin is a yellow-brown pigment, but changes to green due to oxidation induced by Fouchet reaction. Green bilirubin can easily be detected on yellowish and pink colored background.

for 100 tests FVG-100T
3x100 mL FVG-K-100




Giemsa HP kit

Four-reagent kit for staining *Helicobacter pylori* in gastroscopic sections according to Lennart. Advantages of this method for detecting *H. pylori* are sensitive and reproducible results and easy performance.

for 100 tests GMHP-100T






Gomori Trichrome kit

Five-reagent kit for staining muscle, collagen fibre and nuclei, contains blue counterstain. Kit can be used to contrast skeletal, cardiac or smooth muscle.


for 100 tests GT-100T
5x100 mL GT-K-100
* Green counterstain also available



Grimelius kit

Five-reagent kit for staining argyrophilic granules. Grimelius kit can be used for detection of secretory intracytoplasmic granules specific for carcinoid tumors and for identification of neuroendocrine cells.

for 100 tests GRI-100T
5x100 mL GRI-K-100




Grocott kit, stabilized

Seven-reagent kit for visualization of fungi and histological argentaaffin structures in general (such as basal membranes). Green counterstain provides clear and visually rich contrast to target structures stained black.

for 100 tests GRC-100T
300-350 tests GRC-K-100


Stable at room temperature



H.B.F.P. kit

Three-reagent Hematoxylin-Basic Fuchsin-Picric acid staining kit for detection of cardiac muscle changes after ischemia or myocardial infarction. H.B.F.P. kit is a non-enzymatic histochemical technique used for identification of early myocardial ischemia occurring on the cardiac muscle during the first six hours of symptoms.


for 100 tests HBFP-100T
3x100 mL HBFP-K-100



Hematoxylin P.T.A. kit

Four-reagent Hematoxylin-Phosphotungstic Acid kit for differentiation of smooth and striated muscle tissues as well as for detection of fibrin, collagen and elements of central nervous system according to Mallory.

for 100 tests HPTA-100T
4x100 mL HPTA-K-100




Hematoxylin W kit

Acid-resistant hematoxylin according to Weigert. Two-reagent kit that stains the nuclei blue-black, often a component of special staining kits for connective tissues.

for 100 tests HEMW-100T
2x100 mL HEMW-K-100
2x500 mL HEMW-K-500
2x1000 mL HEMW-K-1L






HemoGnost Perls kit

Three-reagent HemoGnost Perls (Prussian blue / Berlin blue) kit for the detection of reactive ferric (Fe³⁺) ions in tissue cells (like spleen or endometrial tissue). When staining bone marrow or other hematology tissue, use LeukoGnost PERLS kit (page 32).


for 100 tests HP-100T
3x250 mL HP-K-250



Ladewig Trichrome kit

Four-reagent kit for staining muscle (brown to red) and collagen fibers with blue counterstain. Staining procedure is easier and faster compared to Masson and Mallory Trichrome staining procedures.


for 100 tests LDW-100T
4x100 mL LDW-K-100



LeukoGnost SPE kit

Kit for detection of specific esterase activity (Leder stain, chloroacetate esterase) on formalin-fixed, paraffin-embedded (FFPE) samples of bone marrow tissue.

for at least 100 tests LKG-SPE



Luxol Fast Blue kit

Three-reagent kit for staining myelin and myelinated axons, Nissl bodies and phospholipids according to Kluver-Barrera. Kit is used for identifying the basic neuronal brain structure or spinal cord sections.


for 100 tests LFB-100T
3x100 mL LFB-K-100



M.I.F. kit

Two-reagent Merthiolate-Iodine-Formalin kit for fixing and staining fecal parasites (especially protozoae, cysts, helminth eggs and larvae). Stool sample is fixed with formalin and stained with two coloring agents; iodine and Eosin Y.


for 100 tests MIF-100T
for 200 tests MIF-200T



Mallory Trichrome kit

Three-reagent staining kit for connective tissue visualization and detection of collagen, cartilage, muscle, elastic fibers, mucous, pituitary cells, reticulum, bones, amyloid and erythrocytes.


for 100 tests MT-100T
3x100 mL MT-K-100



Martius Scarlet Blue (MSB) kit

Seven-reagent kit used for fibrin visualization, especially of older clusters. This method is a modification of Masson Trichrome and is ideal for studying connective tissue and vascular pathology.


for 100 tests MSB-100T
6x100 mL + 1x250 MSB-K-100
mL



Masson Fontana kit

Six-reagent melanine and argentaffin granule staining kit, based on the reduction of silver nitrate to elemental silver. Melanin is a brown-black pigment normally present in hair, skin, retina, iris and certain parts of CNS. Argentaffin granules are found in carcinoid tumors.


for 100 tests MF-100T
9x100 mL MF-K-100
9x500 mL MF-K-500



Masson-Goldner Trichrome kit

Seven-reagent kit for staining muscle and collagen fibers with green counterstain. Also used for visualizing gametes, nuclei, neurofibrils, glial cells, keratins, intercellular fibrils and for differentiation of smooth muscle fibers and collagens.


for 100 tests MGT-100T
7x100 mL MGT-K-100
7x500 mL MGT-K-500



Masson Trichrome kit

Seven-reagent kit for staining muscle and collagen fibers with blue counterstain. Also used for visualizing gametes, nuclei, neurofibrils, glial cells, keratins and intercellular fibrils. Kit may be useful for differentiation between collagen and smooth muscle fibers or detection of diseases like cirrhosis.


for 100 tests MST-100T
7x100 mL MST-K-100
7x500 mL MST-K-500



Movat kit

Eleven-reagent kit for staining collagen, elastic and muscle fibers, mucin and fibrin in tissue section. Movat kit is particularly useful when examining heart and vascular diseases.

for 100 tests MOV-100T
11x100 mL MOV-K-100




Mucicarmine kit

Mucicarmine kit is often used to identify primary tumor sites, distinguishing mucin-negative undifferentiated squamous cell lesions from mucin-positive adenocarcinomas. Can be also used for detection of diseases such as asthma, bronchitis, chronic obstructive pulmonary disease and cystic fibrosis.

for 100 tests MUC-100T
5x100 mL MUC-K-100






Oil Red O kit

Four-reagent kit for selective staining and detection of fat cells and neutral fats according to Johnson. Can be used with frozen sections and fresh smears to detect obesity-linked pathologies such as dyslipidemia and diabetes.


for 100 tests ORO-100T
4x250 mL ORO-K-250



Orcein kit

Five-reagent kit for visualization of hepatitis B surface antigen (HBsAg) seen as viral inclusion bodies in hepatocytes, for elastic fibers and copper associated protein in tissue sections. Can be used with frozen sections.


for 100 tests ORC-100T
5x250 mL ORC-K-250



Paraldehyde Fuchsin kit

Seven-reagent kit according to Gomori for detecting pathological changes in elastic fibers. It also stains mast cell granules, beta granules in pancreatic islets, neurosecretory material, mast cell granules and beta cells in the pituitary gland.


for 100 tests PAF-100T
9x100 mL PAF-K-100



P.A.S. kit

Five-reagent Periodic Acid-Schiff kit for staining aldehydes, mucopolysaccharides, mucoproteins and lymphocytes according to Hotchkiss-McManus. P.A.S. staining may also be used for the demonstration of fungal organisms in tissue sections.


for 100 tests PAS5-100T
5x100 mL PAS5-K-100
5x500 mL PAS5-K-500



P.A.S. Diastase kit

Five-reagent kit for detecting glycogen and mucopolysaccharide structures mostly in liver. The diastase digestion procedure, followed by P.A.S. staining, is useful in the evaluation of glycogen storage disease.

for 100 tests PDIA-100T




P.A.S.M. / Jones kit, stabilized

Seven-reagent Periodic Acid Silver Methenamine kit for staining kidney glomerular basement membranes. Kit includes red counterstain which provides visually rich contrast to target structures stained black.

for 100 tests PASM-100T
300-350 tests PASM-K-100


Stable at room temperature



Perls-Van Gieson kit

Three-reagent kit for the detection of reactive ferric (Fe³⁺) ions in cells (Prussian blue/Berlin blue reaction) with Fuchsin Acid Van Gieson counterstain to differentiate collagen from connective tissue


for 100 tests PEG-100T
3x100 mL PEG-K-100



Picro-Sirius Red kit

Three-reagent kit for collagen staining and visualization in histology. Contains Sirius Red dye which enhances collagen birefringence so it can easily be identified by observing the section under polarized light.


for 100 tests PSR-100T



Reticulin kit

Seven-reagent kit for detecting argyrophilic reticulin fibers. It clearly differentiates between collagen and reticulin and nerve fibers and connective tissue. The main function of reticular fibers is to provide support and are normally found in liver, lymph nodes, spleen and kidneys.


for 100 tests RE-100T
7x50 mL RE-K-50
7x100 mL RE-K-100



Reticulin Contrast kit

Nine-reagent kit for detecting argyrophilic reticulin fibers according to Gordon and Sweets. The kit contains gold chloride solution that enhances visualization of reticulin fibers and it also contains Nuclear Fast Red (Kernechtrot) reagent that enables fine contrasting background.


for 100 tests RET-100T
9x50 mL RET-K-50
9x100 mL RET-K-100



Rhodanine kit

Four-reagent kit for detecting copper and copper-associated protein (CAP) in patients suffering from Wilson's disease. Abnormal copper accumulations are predominately found in liver tissue, but can also be found in the brain and cornea.


for 100 tests ROD-100T
5x100 mL ROD-K-100



Safranin O kit

Four-reagent kit for identification of cartilage, mucins and mastocyte granules used in histology and cytology.


for 100 tests SAFO-100T
4x100 mL SAFO-K-100



Sperm-Diff RTU kit

Ready-to-Use three-reagent kit with reagents stored in containers that can be used as staining jars. Kit contains fixative reagent, red and blue components for fast and effective staining.


3x100 mL SP-RTU-100



Sudan Black B Lipid kit

Four-reagent kit for specific lipid staining in cytochemistry. Sudan Black dye stains a few types of lipids, including neutral fats, phospholipids and sterols. Contains double amount of Sudan Black B, solution.


for 100 tests SBB-100T
6x100 mL SBB-K-100



TB-Stain Histo kit

Three-reagent kit for staining acid-fast bacteria (pathogenic mycobacteria) in histology sections, sputum, smears and culture smears according to Ziehl-Neelsen. Heating of the carbol-fuchsin solution is avoided in this protocol hence the release of hazardous phenolic vapors is omitted.

for 100 tests TBHK-100T



UriGnost S kit

Kit for sampling, staining, and microscopic analysis of urine sediment. Contains modified reagent according to Sternheimer (reagent manufactured according to the European Confederation of Laboratory Medicine (ECLM) and all the necessary equipment for sampling, concentrating, staining, and counting cells and kidney cylinders, as well as urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 µL pipette tips and urine plates).

for 500 tests USK-500

* for more information, see page 59




UriGnost SM kit

Kit for sampling, staining and microscopic analysis of urine sediments. Contains modified reagent according to Sternheimer-Malbin and equipment required for sampling, concentration, staining, counting of kidney cells and kidney cylinders and urine sediment analysis (test tubes with retentive bottom, Eppendorf 200 µL pipette tips and urine plates).

for 500 tests USMK-500

* for more information, see page 59




Van Gieson Trichrome kit

Three-reagent kit for staining collagen fibers, muscle tissue, keratinized epithelium, cytoplasm, glial fibers and erythrocytes. Used for differentiation between collagen and smooth fibers in tumors and various other diseases.

for 100 tests VG-100T
3x100 mL VG-K-100
3x500 mL VG-K-500






Verhoeff kit

Six-reagent kit for detecting atrophy of elastic tissue in cases of emphysema, thinning and loss of elastic fibers in arteriosclerosis and other vascular diseases, or whether blood vessels have been invaded by a tumor.


for 100 tests VER-100T
6x100 mL VER-K-100



Von Kossa kit

Five-reagent kit for simple and reproducible detection of calcium deposits and calcium salt in tissue samples according to Von Kossa. Tissue calcification is associated with metabolic problems in various tissues (like bone marrow and mamma) and in tumors.

for 100 tests VK-100T
5x100 mL VK-K-100



Warthin Starry kit

Five-reagent kit for staining *Spirochaeta spp.*, *Helicobacter pylori*, *Microsporidia spp.* and *Legionella pneumophila*. The kit contains 12 jars with gelatin that enables both incubation and staining of sections, as well as other reagents that enable precipitation of silver on the bacterial surface. The bacteria are found in the mucus of the surface epithelium, in the apical gastric glands and in the gastric mucosa.

up to 48 tests WS-12T



Weigert-Van Gieson kit

Six-reagent kit for staining elastic fibers with prolonged incubation period. Used for differentiation between elastic tissue, collagen and other types of connective tissue.

for 100 tests WVG-100T
6x100 mL WVG-K-100





Additional staining reagents and solutions

Ammonium water

For use with Bielschowsky kit.

100 mL AV-OT-100

Ammonium iron sulfate, solution

For use with Reticulin and Reticulin Contrast staining kits.

100 mL ASF-OT-100

Aniline, alcoholic solution

For use with Azan Trichrome kit.

100 mL ANA-OT-100
250 mL ANA-OT-250

Aniline Blue reagent

Counterstain used with Masson Trichrome kit.

100 mL ABR-OT-100
500 mL ABR-OT-500

Aniline Blue, solution

For use with MSB kit.

100 mL ABO-OT-100

Azan reagent

For use with Azan Trichrome kit.

100 mL AZR-OT-100
250 mL AZR-OT-250

Azocarmine, solution

For use with Azan Trichrome kit.

100 mL AZ-OT-100
250 mL AZ-OT-250

Basic Activation buffer

For use with Oil Red O kit.

250 mL BAP-OT-250

Benedict's reagent

For detecting glucose in urine.

100 mL BNR-OT-100
250 mL BNR-OT-250
500 mL BNR-OT-500
1000 mL BNR-OT-1L

Biebrich Scarlet-Acid Fuchsin reagent

For use with special kits.

100 mL BSAF-OT-100
500 mL BSAF-OT-500

BioSchiff reagent

Schiff's reagent for detecting aldehydes and mucous substances. A component of several special staining kits.

100 mL BS-OT-100
500 mL BS-OT-500
1000 mL BS-OT-1L

BioSchiff Forte

Reagent for use with LeukoGnost PAS kit. Enhanced formulation of Schiff reagents for staining cytoplasm of neutrophils, monocytes, lymphocytes and basophils intensive magenta.

100 mL BSF-OT-100
500 mL BSF-OT-500

Borax, solution

For use with Grocott and P.A.S.M./Jones kits.

100 mL BO-OT-100
105 mL BO-OT-105
250 mL BO-OT-250

SCHIFF'S REAGENT
STABLE AT 22°C*



* In an unopened original packaging. Keep refrigerated after first use.



Additional staining reagents and solutions

Fuchsin Acid Van Gieson reagent

For use in special staining kits.
Synonym: Picrofuchsin reagent.

100 mL FAG-OT-100

Fuchsin Basic reagent

For use in bacteriology.

100 mL FBR-OT-100
500 mL FBR-OT-500
1000 mL FBR-OT-1L

Fuchsin Basic, solution

For use with H.B.F.P. kit.

100 mL FBO-OT-100

Gelatin, gel in transport jar

For preparing developing solution with Warthin Starry staining kit.

12x8 mL GEL-128



Glutaraldehyde, 4% buffered solution

Solution used with Sudan Black B kit.

100 mL G4-OT-100



Gold chloride solutions

Aqueous solutions of gold chloride (aurichloric acid, HAuCl_4), component of many special staining kits.

Gold chloride, 0.2% solution	100 mL	ZK02-OT-100
Gold chloride, 0.6% solution	100 mL	ZK06-OT-100
Gold chloride, 1% solution	50 mL	ZK1-OT-50
Gold chloride, 2% solution	50 mL	ZK2-OT-50

Hematoxylin, Verhoeff A

For use with Verhoeff and Movat kits.

100 mL HEMV-OT-100



Hematoxylin, Weigert A

Hematoxylin solution in ethanol, component of Hematoxylin W kit.
Part of several special staining kits.

100 mL	HEMA-OT-100
500 mL	HEMA-OT-500
1000 mL	HEMA-OT-1L



Hydroquinone, solution

For preparation of developing solution in Warthin Starry kit.

50 mL HQ-OT-50



Hydroquinone-sodium sulphite

For use with Grimelius kit

15 x 3,5 g HNS-15-35

Iodine solution, Verhoeff C

For use with Verhoeff and Movat kits.

100 mL JODV-OT-100

Iron chloride, solution

For use with Fouchet-Van Gieson kit.

100 mL FK-OT-100

Kovac indole reagent

Reagent for identification of indole-positive and indole-negative microorganisms.

100 mL KOV-OT-100



Ladewig's reagent

For use with Ladewig Trichrome kit, contains Aniline Blue, Acid Fuchsin and Orange G dyes.

100 mL LDWR-OT-100

Light Green S.F. reagent

Counterstain used with trichrome special staining kits.

100 mL LGS-OT-100

Lithium carbonate, Luxol

Lithium carbonate solution used with Luxol Fast Blue kit.

100 mL LKL-OT-100

Lithium carbonate, solution

Lithium carbonate solution used with Von-Kossa staining kit.

100 mL LK-OT-100

Lugol's solution, M.I.F.

For use with M.I.F. kit.

100 mL LUGM-OT-100

Luxol Fast Blue, solution

For use with Luxol Fast Blue kit.

100 mL LFB-OT-100



M.F. reagent

For use with M.I.F. kit.

100 mL MF-OT-100

Martius Yellow, solution

For use with MSB kit.

100 mL MAY-OT-100

Mayer's mucicarmine

For use with Mucicarmine kit.

100 mL MMU-OT-100



Metanil Yellow, solution

For use with Mucicarmine kit.

100 mL MY-OT-100



Additional staining reagents and solutions

Potassium hexacyanoferrate, solution

Potassium hexacyanoferrate (II) solution, optimally concentrated for use with HemoGnost Perls kit.

100 mL	KHC-OT-100
250 mL	KHC-OT-250

Potassium hydroxide, solution

Optimally concentrated aqueous solution of potassium hydroxide used with Congo Red Highman kit.

100 mL	KHO-OT-100
--------	------------

Potassium permanganate, solutions

For use with Hematoxylin P.T.A., Masson Fontana, Orcein, Paraldehyde Fuchsin and Reticulin staining kits.

Potassium permanganate, 0.5% solution	100 mL	KP05-OT-100
---------------------------------------	--------	-------------

Potassium permanganate, 1% solution	250 mL	KP1-OT-250
-------------------------------------	--------	------------

Propylene glycol

For use with Sudan Black B Lipid kit.

100 mL	PG-100
--------	--------

Propylene glycol, 85%

For use with Sudan Black B Lipid kit.

100 mL	PG85-100
--------	----------

P.T.A. - P.M.A. reagent

Reagent containing optimal concentration of phosphotungstic and phosphomolybdic acid. Used with Masson Trichrome and Masson-Goldner Trichrome staining kits.

100 mL	PPR-OT-100
500 mL	PPR-OT-500

Reagent for differentiation in Verhoeff kit

For use with Verhoeff kit.

100 mL	RDV-OT-100
--------	------------

Reducing buffer, solution

For use with Von Kossa kit.

100 mL	RP-OT-100
--------	-----------

Resorcine Fuchsin reagent

For use with Weigert-Van Gieson kit.

100 mL	RFR-OT-100
--------	------------

Rhodanine reagent

For use with Rhodanine kit.

100 mL	RR-OT-100
--------	-----------

Safranin O reagent

Counterstaining reagent for use in histology and cytology. Used for visualizing and staining cartilage, mucins and mastocytes in formalin-fixed and paraffin-embedded tissue samples.

100 mL	SAFO-OT-100
--------	-------------

Silver nitrate solutions

For use in special staining kits.

Silver ammonia solution (component of Reticulin and Reticulin Contrast staining kits) ●	100 mL	SA-OT-100
---	--------	-----------

Silver ammonia reagent (component of Von Kossa and Masson Fontana staining kits) ●	100 mL	SAR-OT-100
--	--------	------------

Silver ammonia reagent stock solution (component of Bielschowsky staining kit) ●	100 mL	SSAR-OT-100
--	--------	-------------

Silver nitrate, 20% solution (component of Bielschowsky staining kit) ●	100 mL	SN20-OT-100
---	--------	-------------

Silver nitrate solution, stabilized (component of Grocott and P.A.S.M. / Jones, stabilized staining kits)	100 mL	SNS-OT-100
---	--------	------------

Silver nitrate, WS solution (component of Warthin Starry staining kit) ●	100 mL	SNWS-OT-100
--	--------	-------------

Sodium acetate, solution

For use with Rhodanine kit.

100 mL	NA-OT-100
--------	-----------

Sodium hydroxide, solution

For use with Congo Red Puchtler kit.

30 mL	NH-OT-30
-------	----------

Sodium metabisulfite solutions

Component of Alcian Blue-P.A.S., P.A.S. and Feulgen staining kits.

Sodium metabisulfite	100 mL	NM-OT-100
	500 mL	NM-OT-500

Sodium metabisulfite, solution for Feulgen	100 mL	NMF-OT-100
--	--------	------------

Sodium tetraborate, solution

A component of Alcian Blue pH 2.5 and Alcian Blue-P.A.S. staining kits.

100 mL	NTB-OT-100
500 mL	NTB-OT-500

Sodium thiosulfate solution

Solutions are component of Grocott stabilized, Masson Fontana, P.A.S.M. / Jones, stabilized, Reticulin, Reticulin Contrast and Von Kossa staining kits.

Sodium thiosulfate, 2% solution	100 mL	NT2-OT-100
---------------------------------	--------	------------

Sodium thiosulfate, 5% solution	100 mL	NT5-OT-100
---------------------------------	--------	------------

Sudan Black B, solution

Solution for staining lipids, used with Sudan Black B lipid kit.

100 mL	SBO-OT-100
--------	------------



Hematoxylin, C.I. 75290

Natural Black 1, BSC certified stain. For preparing nuclear staining reagents.

25 g	H-P-25
100 g	H-P-100

Janus Green B, C.I. 11050

Diazine Green, BSC certified stain. For displaying mitochondria in living cells.

25 g	JG-P-25
------	---------

Jenner Stain

Jenner, BSC certified stain. Romanowsky polychromatic stain routinely used in hematology for staining blood smears.

25 g	JS-P-25
------	---------

Leishman Stain

Eosin-polychrome Methylene Blue. Romanowsky polychromatic dye. Used to distinguish and identify leukocytes and malaria parasites.

25 g	LS-P-25
------	---------

Light Green S.F., C.I. 42095

Acid Green 5, Pencil Green SF, BSC certified stain. For counterstaining according to Papanicolaou and Masson trichrome methods.

25 g	LG-P-25
------	---------

Luxol Fast Blue MBSN, C.I. 74180

Luxol Fast Blue, Solvent Blue 38. For use in histology.

25 g	LFB-P-25
------	----------

Malachite Green, C.I. 42000

Malachite Green Oxalate, Basic Green 4, Victoria Green, BSC certified stain. Used as counterstain in TB-Stain Cold kit.

10 g	MAG-P-10
25 g	MAG-P-25

Martius Yellow C.I. 10315

Golden Yellow, Martinsgelb, Acid yellow 24. For staining erythrocytes yellow in trichrome staining methods.

25 g	MAY-P-25
------	----------

May-Gruenwald

Polychromatic Romanowsky dye. For blood smears and bone marrow staining.

25 g	MG-P-25
50 g	MG-P-50

Metanil Yellow, C.I. 13065

Acid yellow 36. For staining cytoplasm, muscle connectors and identifying parietal cells of gastric glands.

25 g	MEY-P-25
50 g	MEY-P-50
100 g	MEY-P-100

Methyl Green, C.I. 42590

For staining nuclei green.

25 g	MGR-P-25
------	----------

Methyl Violet 2B, C.I. 42535

Basic Violet 1, BSC certified stain. For bacterial staining according to Gram.

25 g	MV2B-P-25
------	-----------

Methyl Violet 10B, C.I. 42555

Gentian Violet, Crystal Violet, BSC certified stain. For bacterial staining according to Gram.

25 g	MV10B-P-25
50 g	MV10B-P-50

Methylene Blue, C.I. 52015

Basic Blue 9, BSC certified stain. Used for preparing Romanowsky polychromatic stain for staining bone marrow and as a counterstain in staining of acid-fast bacteria.

25 g	MB-P-25
100 g	MB-P-100

Methylene Green, C.I. 52020

Methylene Green, BSC certified stain. For staining nuclei green.

25 g	MEG-P-25
------	----------

Neutral Red, C.I. 50040

Basic Red 5, BSC certified stain. For supravital staining of blood cells using Neutral Red-Janus Green B method.

25 g	NR-P-25
------	---------

Nigrosine WS, C.I. 50420

Nigrosine water soluble, Nigrosine B. For use in microbiology and cytology, as negative staining.

25 g	NGR-P-25
------	----------

Nuclear Fast Red, C.I. 60760

Kernechtrot. Used for staining nuclei red by creating aluminum complex.

5 g	NF-P-5
-----	--------

Oil Red O, C.I. 26125

Sudan Red 5B, BSC certified stain. For staining lipids.

25 g	ORO-P-25
------	----------

Orange II, C.I. 15510

Acid Orange 7, BSC certified stain. Used as a counterstain, as a substitute for Orange G.

25 g	O2-P-25
------	---------

Orange G, C.I. 16230

Acid Orange 10, Acid Fast Orange G, BSC certified stain. For staining keratin according to Papanicolaou, counterstain for hematoxylin.

25 g	OG-P-25
100 g	OG-P-100

Orcein, C.I. (ed.1) 1242

Natural Red 28, Orcein Synthetic. For staining elastic tissues and viral inclusion bodies of the hepatitis B surface antigen.

5 g	ORC-P-5
10 g	ORC-P-10



Polypropylene dissecting boards

Boards for pathohistological tissue dissection made of polypropylene.

Polypropylene, white, dimensions 60x40x1.5 cm	1 pc	BC001
Polypropylene, dark blue, dimensions 39x29.5x2.5 cm	1 pc	BC032
Polypropylene, white, dimensions 40x30x1.5 cm	1 pc	BC061CS

Bio/Cut microtome blades - high profile

Bio/Cut H35C microtome blades for cryotome, high profile (wide). Ideal for frozen sections. Suitable for both soft and hard tissues.

50 pcs	Bio/Cut H35C
--------	--------------

Bio/Cut microtome blades - low profile

Bio/Cut 35A microtome blades for soft and hard tissues, low profile.

50 pcs	Bio/Cut 35A
--------	-------------



Polypropylene containers for sample transport

Container with cover for transporting histology samples and organs.

500 mL	1 pc	PK500
1000 mL	1 pc	PK1000
1200 mL	1 pc	PK1200
2500 mL	1 pc	PK2500
3750 mL	1 pc	PK3750
5000 mL	1 pc	PK5000
7000 mL	1 pc	PK7000
10000 mL	1 pc	PK10000



Microtome oil

Optimal viscosity oil for microtome maintenance.

Microtome oil 50 mL	1 pc	MO-50
Microtome oil 100 mL	1 pc	MO-100



Contact and ordering information

BioGnost Ltd.,
Medjugorska 59, Zagreb
10040 Croatia (Europe)

WORKING HOURS
Monday through Friday
8:00 to 16:00 (CET)

e-mail: sales@biognost.hr
website: www.biognost.com
phone: +385 1 2409 997
fax: + 385 1 2404 039



Technical support

Technical questions can be answered by emailing our sales representatives at sandra.ban@biognost.hr and viktorija.liber@biognost.hr.

MSDS and product information are available on our website at www.biognost.com.

Frequently Asked Questions (FAQ) section is available on our website.



Shipping information

Shipping and handling charges are prepaid and added to the invoice. If requested, we can use your account number or carrier of your choice, and also ship on your third-party freight accounts (Incoterms 2020). Charges vary according to destination, weight and content, and are available upon request.

Our employees are trained to classify, package, mark and label products identified as dangerous goods according to ADR, IMDG and IATA to ensure safe transport.



Index

4% Formaldehyde	8
4% Formaldehyde, alcoholic solution	53
4% NB Formaldehyde	8
4% NB Formaldehyde in transport containers	9
10% Formaldehyde	9
10% NB Formaldehyde	9

A

Acetic acid, 0.5% solution	55
Acetic acid, 1% solution	55
Acetic acid, Colloidal Iron (A) solution	55
Acetic acid, Colloidal Iron (B) solution	55
Acetic buffer, Grimelius	53
Acid alcohol	26
Acid alcohols	53
Acid solution	53
Acridine Orange, C.I. 46005	60
Acridine Orange solution	53
A.F.O.G. kit	43
A.F.O.G. reagent	53
Alcian Blue 8 GX, C.I. 74240	60
Alcian Blue - P.A.S. kit	43
Alcian Blue pH 1.0 kit	43
Alcian Blue pH 2.5 kit	43
Alcian Blue solution 1%	53
Alcian Blue solution pH 1.0	53
Alcian Blue solution pH 2.5	53
Alcian Green	60
Alcian Yellow, C.I. 12840	60
Alcian Yellow Toluidine Blue kit	43
Alkaline alcohol, solution	53
Amido Black 10B, C.I. 20470	60
Ammonium iron sulfate, solution	54
Ammonium water	54
Aniline, alcoholic solution	54
Aniline Blue, C.I. 42780	60
Aniline Blue reagent	54
Aniline Blue, solution	54
Auramine O, C.I. 41000	60
Azan reagent	54
Azan Trichrome kit	43
Azocarmine G, C.I. 50085	60
Azocarmine, solution	54
Azophloxine, C.I. 18050	60
Azure A, C.I. 52005	60
Azure B, C.I. 52010	60
Azure C, C.I. 52002	60
Azure II	60
Azure II Eosinate	61

B

Basic Activation buffer	54
Benedict's reagent	54
Biebrich Scarlet-Acid Fuchsin reagent	54
Biebrich Scarlet, C.I. 26905	61
Bielschowsky kit	44
BioBluing buffer	26
BioClear	14
BioClear New	14
Bio/Cut microtome blades - high profile	65
Bio/Cut microtome blades - low profile	65
Bio-Diff 1 reagent	30
Bio-Diff 1 RTU reagent	31
Bio-Diff 2 reagent	30
Bio-Diff 2 RTU reagent	31
Bio-Diff 3 reagent	30
Bio-Diff 3 RTU reagent	31
Bio-Diff kit	30
Bio-Diff RTU kit	31, 44
BioFix	10
BioFix C	10

BioFix GL	10
BioFix GL in transport containers	10
BioFix HD	10
BioForm S	10
BioGram 4 kit	36
BioGram ECO kit	37
BioGram Histo kit	44
Bio-Hansel solution	33
BioMount	19
BioMount Aqua	20, 35
BioMount C	20
BioMount DPX	18
BioMount DPX High	18
BioMount DPX Low	18
BioMount DPX New	18
BioMount High	19
BioMount M	19
BioMount New	19
BioMount New Low	19
Biopsy pads for cassettes, blue	64
BioSchiff Forte	54
BioSchiff reagent	54
BioWax 52/54	15
BioWax 56/58	15
BioWax Blue	15
BioWax Plus 56/58	15
Bismarck Brown Y, C.I. 21000	61
Bluing reagent	26
Borax, solution	54
Bouin's solution	11
Brilliant Cresyl Blue, C.I. 51010	61
Brilliant Cresyl Blue, solution	33
Brilliant Green Certified, C.I. 42040	61
Buffer solution, pH 6.4	32
Buffer solution, pH 6.8	32
Buffer solution, pH 7.2	31
Buffer tablets pH 6.8	32
Buffer tablets pH 7.2	31

C

Carbol Fuchsin Gram 10x concentrated	37
Carmine, C.I. 75470	61
Carminic acid, C.I. 75470	61
Celestine Blue, C.I. 51050	61
Chromotrope 2R / Aniline Blue reagent	55
Chromotrope 2R, C.I. 16570	61
Chromotrope 2R / Fast Green F.C.F. reagent	55
CitoFix	11
CitoSpray	11
CitoSpray solution	11
Clean Lens, solution 1	22
Clean Lens, solution 2	22
Clean Lens, solution 3	22
Colloidal Iron kit	44
Colloidal Iron, stock solution	55
Congo Red, C.I. 22120	61
Congo Red Highman kit	44
Congo Red Puchtler kit	44
Congo Red Puchtler reagent	55
Congo Red reagent	55
Cresyl Violet Acetate Certified	61
Cresyl Violet, solution	55
CryoFix Gel	16
CryoFix Gel blue	16
CryoFix Gel green	16
CryoFix Gel pink	16
CryoFix Gel yellow	16
Cryo Spray	16
Crystal Violet, C.I. 42555	61
CytoGnost kit	33

D

DeHematin	9
DeMold solution	16
DeMold Spray	16

L

Lactophenol Blue solution	41
Lactophenol solution	41
Ladewig's reagent	56
Ladewig Trichrome kit	47
Leishman's solution	33
Leishman Stain	62
LeukoGnost ACP	34
LeukoGnost ALP	34
LeukoGnost Fixative	35
LeukoGnost Fixative ECO	35
LeukoGnost Fixative GL	35
LeukoGnost HEM	35
LeukoGnost MPO	34
LeukoGnost NSE	34
LeukoGnost PAS	34
LeukoGnost PERLS	34
LeukoGnost PLUS	35
LeukoGnost SPE	34
LeukoGnost SPE kit	47
LeukoGnost SPENSE	34
Lid for cassettes	64
Light Green S.F., C.I. 42095	62
Light Green S.F. reagent	56
Liquid Blocker (PAP pen)	7
Lithium carbonate, Luxol	56
Lithium carbonate, solution	56
Lugol's C solution	37
Lugol's solution	37
Lugol's solution, acid	37
Lugol's solution, M.I.F.	56
Lugol's solution (Schiller's test)	37
Luxol Fast Blue kit	47
Luxol Fast Blue MBSN, C.I. 74180	62
Luxol Fast Blue, solution	56

M

Malachite Green, C.I. 42000	62
Mallory Trichrome kit	47
Martius Scarlet Blue (MSB) kit	48
Martius Yellow C.I. 10315	62
Martius Yellow, solution	56
Masson Fontana kit	48
Masson-Goldner Trichrome kit	48
Masson Trichrome kit	48
Mayer's mucicarmine	56
May-Gruenwald	62
May-Gruenwald solution	32
Metal histology mold	64
Metanil Yellow, C.I. 13065	62
Metanil Yellow, solution	56
Methenamine, solution	57
Methyl Blue Eosin, solution	57
Methylene Blue 10x concentrated	38
Methylene Blue, C.I. 52015	62
Methylene Blue Loeffler's reagent	38
Methylene Green, C.I. 52020	62
Methyl Green, C.I. 42590	62
Methyl Violet 2B, C.I. 42535	62
Methyl Violet 10B, C.I. 42555	62
M.F. reagent	56
Micro Clean	15
Microtome oil	65
M.I.F. kit	47
Movat kit	48
Mucicarmine kit	48
Myco-Mount Blue	41
Myco-Mount Red	41

N

NB Formaldehyde concentrated 5x	9
Neutral Red, C.I. 50040	62

Neutral Red reagent	57
Nigrosine, isotonic solution	57
Nigrosine WS, C.I. 50420	62
Nuclear Fast Red, C.I. 60760	62
Nuclear Fast Red (Kernechtrot) reagent	57

O

OG-6 reagent, Pap 2A	28
OG-EA31 Pap reagent	29
OG-EA50 Pap reagent	29
Oil Red O, C.I. 26125	62
Oil Red O kit	49
Oil Red O reagent	57
Orange G, 1% solution	57
Orange G / Aniline Blue reagent	57
Orange G, C.I. 16230	62
Orange II, C.I. 15510	62
Orange II reagent, Pap 2B	28
Orcein, C.I. (ed.1) 1242	62
Orcein kit	49
Orcein reagent	57
OsteoFast 1	12
OsteoFast 2	12
OsteoSens	12
Oxalic acid, 1% solution	57
Oxalic acid, 2% solution	57

P

PAP PEN RED	7
Paraldehyde Fuchsin kit	49
Paraldehyde Fuchsin reagent	57
P.A.S. Diastase kit	49
P.A.S. kit	49
P.A.S.M. / Jones kit, stabilized	49
Periodic acid, 0.5% solution	57
Periodic acid, 0.8% solution	57
Periodic acid, 1% solution	57
Perls-Van Gieson kit	50
Phloxine B, C.I. 45410	63
Phosphomolybdic acid, 1% solution	57
Phosphomolybdic acid, 5% solution	57
Phosphotungstic acid, 1% solution	57
Phosphotungstic acid, 5% solution	57
Picric acid in acetone, solution	57
Picro-Sirius Red kit	50
Picro-Sirius Red reagent	57
Polypropylene containers for sample transport	65
Polypropylene dissecting boards	65
Ponceau S, C.I. 27195	63
Ponceau S, solution	57
Potassium hexacyanoferrate, Colloidal Iron solution	57
Potassium hexacyanoferrate, solution	58
Potassium hydroxide, solution	58
Potassium hydroxide, solution for fungi	41
Potassium permanganate	63
Potassium permanganate, 0.5% solution	58
Potassium permanganate, 1% solution	58
Propylene glycol	58
Propylene glycol, 85%	58
P.T.A. - P.M.A. reagent	58
Pyronine B, C.I. 45010	63

R

Rapid Pap staining kit	29
Reagent for differentiation in Gomori Trichrome kit	55
Reagent for differentiation in Verhoeff kit	58
Reducing buffer, solution	58
Resorcine Fuchsin reagent	58
Resorcinol, C.I. 76505	63
Reticulin Contrast kit	50
Reticulin kit	50
Rhodamine B, C.I. 45170	63

Rhodanine	63
Rhodanine kit	50
Rhodanine reagent	58
Rose Bengal, C.I. 45440	63

S

Saccomanno's fixative	11
Safranin O, C.I. 50240	63
Safranin O kit	50
Safranin O reagent	58
Scott's solution	26
Shorr's solution	29
Silver ammonia reagent	58
Silver ammonia reagent stock solution	58
Silver ammonia solution	58
Silver nitrate, 20% solution	58
Silver nitrate solution, stabilized	58
Silver nitrate, WS solution	58
Sirius Red, C.I. 35780	63
Sodium acetate, solution	58
Sodium hydroxide, solution	58
Sodium metabisulfite solutions	58
Sodium tetraborate, solution	58
Sodium thiosulfate, 2% solution	58
Sodium thiosulfate, 5% solution	58
Sodium thiosulfate solution	58
Sperm-Diff RTU kit	51
Sudan Black B, C.I. 26150	63
Sudan Black B ECO	34
Sudan Black B kit	34
Sudan Black B Lipid kit	51
Sudan Black B reagent	59
Sudan Black B, solution	58
Sudan III, C.I. 26100	63
Sudan IV, C.I. 26105	63
Sulfuric acid, 0.3% solution	59
Sulfuric acid, 0.5% solution	59
Sulfuric acid, 3% solution	59
Sulphosalicylic acid, 20% solution	59

T

TB Armand reagent	39
TB Auramine O reagent	40
TB Auramine-Rhodamine reagent	40
TB Carbol Fuchsin reagent	38, 39
TB Decolorizer	38, 39
TB Decolorizer Fluorescent	40
TB-Fuchsin reagent	38
TB Malachite Green reagent	39
TB Permanganate reagent	40
TB-Stain Auramine O kit	40
TB-Stain Cold kit	39
TB-Stain ECO kit	38
TB-Stain Fluorescent kit	40
TB-Stain Histo kit	51
TB-Stain Hot kit	38
TB-Stain Quick kit	39
ThromboGnost solution	33
Tissue marking dye, black	7
Tissue marking dye, blue	7
Tissue marking dye, green	7
Tissue marking dye, lilac	7
Tissue marking dye, orange	7
Tissue marking dye, red	7
Tissue marking dyes, 5 colour set	7
Tissue marking dyes, 7 colour set	7
Tissue marking dye, yellow	7
Tissue Soft	16
Toluidine Blue O, C.I. 52040	63
Toluidine Blue, solution	59
Trichloroacetic acid, solution	59
Tuerk's solution	33

U

UriGnost S kit	51
UriGnost SM kit	51
UriGnost SM reagent	59
UriGnost S reagent	59

V

Van Gieson Trichrome kit	51
Verhoeff kit	52
Victoria Blue B, C.I. 44045	63
Victoria Blue R, C.I. 44040	63
VitroGnost Plus Ultra adhesive microscope slides	64
Von Kossa kit	52

W

Warthin Starry kit	52
Water Blue, C.I. 42755	63
Weigert-Van Gieson kit	52
Wright-Giemsa solution	33
Wright-Giemsa, solution for reptiles	33
Wright's solution	33
Wright's Stain	63

X

Xylidine Ponceau, C.I. 16150	63
------------------------------	----

Be a part of
our global network
of satisfied customers

BioGnost Ltd.

For development, production
and marketing of medical products

Medjugorska 59, 10040 Zagreb, Croatia, EU
phone: +385 1 2409 997, fax: +385 1 2404 039
sales@biognost.hr • www.biognost.com

